


DUDLEY KNOX LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIF 93940

NAVAL POSTGRADUATE SCHOOL

Monterey, California



THESIS

RELATIONSHIPS BETWEEN SEA SURFACE TEMPERATURE
AND NUTRIENTS IN SATELLITE DETECTED OCEANIC
FRONTS

by

John Woepfel Conrad

March 1980

Thesis Advisor:

Eugene D. Traganza

Approved for public release; distribution unlimited

T195209

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Relationships Between Sea Surface Temperature and Nutrients in Satellite Detected Oceanic Fronts		5. TYPE OF REPORT & PERIOD COVERED Master's Thesis; March 1980
7. AUTHOR(s) John Woepfel Conrad		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Naval Postgraduate School Monterey, California 93940		12. REPORT DATE March 1980
		13. NUMBER OF PAGES 114
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nutrients, Nitrate, Phosphate, satellite infrared imagery, thermal fronts, chemical fronts, fronts, Nitrogen, Phosphorus, sea surface temperatures.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Satellite IR images of the California coast off Point Sur reveal recurrent surface features which appear to be "thermal dis- continuities" associated with aperiodic upwelling events. Some of these have associated "chemical fronts" and increased biological activity. Satellite IR imagery was used to locate "discontinuities" and with <u>in situ</u> monitoring the development of three features were studied. Interrelationships between sea surface temperature		

20. (cont'd)

nutrients and microplanktonic biomass were investigated. Nutrient ratios, satellite imagery, wind stress data and correlations between nutrients and temperature were used to develop an estimate of "age" within a simplified upwelling "life cycle" model.

The features range in scale from tens to hundreds of kilometers. Two upwelling features exhibited very strong correlations between nutrient and temperature but a third feature had considerable nutrient variability. This suggests a considerable impact from the dynamic and biological processes. The technique of coupling satellite imagery and in situ monitoring was found to be a feasible method to provide real time inferences of the nutrient structure associated with an upwelled thermal feature.

Approved for public release; distribution unlimited

Relationships Between Sea Surface Temperature
and Nutrients in Satellite Detected
Oceanic Fronts

by

John Woeppel Conrad
Lieutenant Commander, United States Navy
B.S., United States Naval Academy, 1969

Submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE IN OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL
March 1980

ABSTRACT

Satellite IR images of the California coast off Point Sur reveal recurrent surface features which appear to be "thermal discontinuities" associated with aperiodic upwelling events. Some of these have associated "chemical fronts" and increased biological activity. Satellite IR imagery was used to locate "discontinuities" and with in situ monitoring the development of three features were studied. Interrelationships between sea surface temperature, nutrients and microplanktonic biomass were investigated. Nutrient ratios, satellite imagery, wind stress data and correlations between nutrients and temperature were used to develop an estimate of "age" within a simplified upwelling "life cycle" model.

The features range in scale from tens to hundreds of kilometers. Two upwelling features exhibited very strong correlations between nutrient and temperature but a third feature had considerable nutrient variability. This suggests a considerable impact from the dynamic and biological processes. The technique of coupling satellite imagery and in situ monitoring was found to be a feasible method to provide real time inferences of the nutrient structure associated with an upwelled thermal feature.

TABLE OF CONTENTS

I.	INTRODUCTION-----	11
II.	THEORY-----	13
	A. UPWELLING-----	13
	B. SATELLITE IMAGERY-----	14
	C. NUTRIENTS (NITROGEN AND PHOSPHORUS)-----	15
III.	METHODS-----	17
	A. NUTRIENTS-----	17
	B. SATELLITE IMAGERY-----	17
	C. COMPUTATIONS-----	18
	D. CHLOROPHYLL-----	19
	E. ATP-----	19
	F. TEMPERATURE-----	19
	G. WIND-----	20
IV.	RESULTS-----	21
	A. 30 APRIL CRUISE-----	21
	B. 13 JUNE CRUISE-----	32
	C. 7 AUGUST CRUISE-----	33
V.	DISCUSSION-----	54
VI.	CONCLUSIONS-----	62
APPENDIX A.	Listing of Cruise Data: Time, Latitude, Longitude, Elapsed Distance, ATP, Nitrate, Phosphate, Chlorophyll, Nutrient Ratio, Temperature-----	64
APPENDIX B:	Wind Data: Wind Stress, Ekman Transport, Up- Welling Index, Vertical Velocity-----	102
	BIBLIOGRAPHY-----	108
	INITIAL DISTRIBUTION LIST-----	112

LIST OF PHOTOGRAPHIC PLATES

Plate 1.	TIROS-N Satellite IR Image for 19 April 1979----	22
Plate 2.	TIROS-N Satellite IR Image for 29 April 1979----	23
Plate 3.	TIROS-N Satellite IR Image for 13 June 1979-----	34
Plate 4.	TIROS-N Satellite IR Image for 30 July 1979-----	42
Plate 5.	TIROS-N Satellite IR Image for 5 August 1979----	43

LIST OF FIGURES

Fig. 1.	Track of the 30 April Cruise and Outline of Upwelling Feature-----	24
Fig. 2.	Nitrate, Phosphate and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	25
Fig. 3.	ATP, Chlorophyll <u>a</u> , and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	26
Fig. 4.	Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	27
Fig. 5.	Nitrate Versus Phosphate for the 30 April Cruise-	28
Fig. 6.	Nitrate Versus Temperature for the 30 April Cruise-----	29
Fig. 7.	Phosphate Versus Temperature for the 30 April Cruise-----	30
Fig. 8.	Vertical Temperature Sections Along the Track of the 30 April Cruise-----	31
Fig. 9.	Track of the 13 June Cruise and Outline of the Upwelling Feature-----	35
Fig. 10.	Nitrate, Phosphate, Nutrient Ratio, ATP, Chlorophyll <u>a</u> and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 13 June Cruise-----	36
Fig. 11.	Nitrate Versus Phosphate for the 13 June Cruise--	37
Fig. 12.	Nitrate Versus Temperature for the 13 June Cruise-----	38
Fig. 13.	Phosphate Versus Temperature for the 13 June Cruise-----	39
Fig. 14.	Track of the 7 August Cruise and Outline of the Upwelling Feature-----	44

Fig. 15.	Nitrate, Phosphate and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	45
Fig. 16.	ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	46
Fig. 17.	Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	47
Fig. 18.	Nitrate Versus Phosphate for the 7 August Cruise-	48
Fig. 19.	Nitrate Versus Temperature for the 7 August Cruise-----	49
Fig. 20.	Phosphate Versus Temperature for the 7 August Cruise-----	50
Fig. 21.	Vertical Temperature Sections Along the Track of the 7 August Cruise-----	51

LIST OF TABLES

Table I.	Summary of Regression Analyses-----	52
Table II.	Data Ranges-----	53

ACKNOWLEDGMENTS

This thesis is a result of ongoing research in chemical oceanography at the Naval Postgraduate School which is supported by the Office of Naval Research, Code 482, NSTL, Bay St. Louis, Mississippi. I thank the sponsors and key individuals whose assistance made this task possible, including: Dr. Eugene D. Traganza, my thesis advisor and principal investigator; Dr. Christopher N. K. Mooers, Dr. Stevens P. Tucker, Dr. Eugene C. Haderlie, Ms. Andrea McDonald of the Naval Postgraduate School, Monterey, California; Captain W. W. Reynolds and the crew of the R/V ACANIA; Mr. L. Breaker of the National Environmental Satellite Service at Redwood City, California; and Mr. A. Bakun of the National Marine Fisheries Service, Pacific Environmental Group, Monterey, California. Lastly, I thank my family for their forbearance, patience and support.

I. INTRODUCTION

In this thesis satellite thermal imagery and in situ automated biochemical analyses were combined to study inter-relationships of sea surface temperatures, nitrate, phosphate and biological activity in oceanic fronts.

An oceanic front is a region of transition between two oceanic regimes with different characteristics (Cheney, 1976). These characteristics can be temperature, salinity, chemical or biological quantities.

In situ automated biochemical and thermal analyses of surface waters were previously accomplished by others, particularly Kelly, et al. (1975). However, they were not linked to satellite infrared imagery. Due to the marginal quality of satellite-derived results, oceanographers have largely either ignored them or have not been convinced that they could be integrated usefully with classical oceanographic observations (Legeckis, 1978). More sophisticated satellites and sensors, such as the TIROS (Television Infrared Observation Satellites) series and the Advanced Very High Resolution Radiometer (AVHRR) have made it possible to make useful inferences in areas of the ocean that are classically associated with sea surface temperature anomalies. The region of interest in this study is off the central California coast, where "oceanic fronts" and "eddies" appear to form in response to wind driven pulses of cold upwelled

coastal waters. The existence of "chemical fronts" in association with these features was postulated in 1978 by Traganza (1978) and demonstrated in 1978 by Traganza et al. (1980).

Continued study of these frontal systems may be particularly relevant to the interests of the Navy in view of their potential effect on the propagation of sound energy and the capability of sensors to distinguish significant sound signals from the background noises in the ocean. Sound velocity changes across thermal fronts, coupled with changes in nutrient concentrations that lead to increased biological activity, may degrade both active and passive sonar performance through changes in sound propagation loss, increased biological reverberation levels and background noise levels.

.

II. THEORY

A. UPWELLING

Coastal upwelling is caused by the divergence of surface water away from the coast. Cold water rises from subsurface layers to replace this water. When the pycnocline surfaces, relatively nutrient-rich water (which has been trapped below the pycnocline due to density differences) will upwell and enrich the surface waters. In the California coastal area this water is generally derived from a depth no greater than 200 meters offshore (Fairbridge, 1966; Sverdrup, et al., 1942). In the Northern Hemisphere coastal upwelling can be caused by the northerly wind stress prevailing in spring and summer along the western continental coasts. These prevailing winds may vary seasonally off the California coast as the North Pacific subtropical high moves north during the spring and summer, then south during the autumn and winter (Bakun, 1973). Upwelling can also be caused by currents impinging on land masses (Fairbridge, 1966) or winds blowing directly away from shore.

The circulation patterns vary with the width and form of the continental shelf and slope, the wind speed, duration, fetch and latitude. The characteristics of upwelled water may vary depending on the depth from which the water came, the properties of the source water, the characteristic vertical velocity, and the residence time of the upwelled

water in the euphotic zone (SCOR Working Group, 1974). The proximity of the heads of underwater canyons enhances the intensity of the upwelling, (SCOR Working Group, 1974; Codispoti, 1977; Treguer, 1977). Therefore, upwelling can vary from region to region and season to season (Codispoti et al., 1979; Barton et al., 1977).

B. SATELLITE IMAGERY

Infrared (IR) images of the sea surface temperatures were received from the TIROS-N series satellites using an Advanced Very High Resolution Radiometer (AVHRR). The measured quantity is the sea surface radiance which is converted to an equivalent blackbody temperature. Correction factors are applied to account for atmospheric attenuation and radiation from atmospheric gases.

The infrared detectors have a temperature response from approximately -113° to $+47^{\circ}\text{C}$ (Brower et al., 1976). This is displayed in gray tones on photographic film, where the darker shades represent higher temperatures. Since the ocean temperature range is considerably smaller than the detector's range it is possible, through the technique of "image enhancement," to assign the available gray shades to this narrower range. This allows finer resolution of the gray shades that represent the oceanic temperature structure and will better distinguish thermal features.

Cloud cover is often difficult to distinguish in infrared images. In this case a comparison between the visual

the ratio of nitrate to phosphate is not very often 16:1 in ocean surface waters (Banse, 1974).

III. METHODS

A. NUTRIENTS

Surface nutrient concentrations were determined approximately every 0.6 km except for the 13 June Cruise Leg 3 when engine problems caused speed adjustments. Seawater samples were pumped from a keel intake at 2.5m to the ship-board laboratory. Nitrate and phosphate were analyzed every two minutes according to the Technicon Autoanalyzer Industrial Method 175-72-WM, 177-72-WM (Anonymous, 1973) and 100-70-WM (Anonymous, 1978). Here nitrate may include traces of in situ nitrite, since the nitrate is reduced to nitrite before measurements. However, according to Paulson (1972) there is little or no interference from surface nitrate in this area.

B. SATELLITE IMAGERY

The pre-cruise procedure was initiated approximately 2-3 weeks prior to the estimated departure time. Mr. Breaker of the National Environmental Satellite Service (NESS) at Redwood City, California, monitored and enhanced images from the TIROS-N satellite series. Several days prior to the cruise, the day of the cruise and sometimes during the cruise updated satellite information was passed via telephone. Information, re IR inferred thermal features, such as orientation, approximate center, size, and spatial relationship

relative to prominent landmarks proved invaluable in locating features and planning sampling strategy. Additionally, the direction of a feature's general movement and large-scale changes in shape may be inferred by viewing a time sequence of images.

C. COMPUTATIONS

A linear regression analysis was performed and mean values were calculated for nitrate, phosphate and temperature using the NORLSQ library subroutine at the W. R. Church Computer Center of the Naval Postgraduate School.

Correlation coefficients for nitrate to phosphate, nitrate to temperature and phosphate to temperature were obtained utilizing the correlation coefficient equation:

$$r = \Sigma (X_i - \bar{X})(Y_i - \bar{Y}) / ([\Sigma (X_i - \bar{X})^2] [\Sigma (Y_i - \bar{Y})^2])^{1/2}$$

Zero values of nitrate and phosphate occurred when taking reference baselines or calibrating standards, or when a nitrate "none detected" condition occurred due to concentrations below the sensitivity of the instrument. When grossly anomalous values occurred, the nutrient and temperature data for that time increment were not used in the calculations.

(Statistical computations for the wind stress, adenosine triphosphate (ATP), and chlorophyll biomass will be discussed in a related thesis.)

D. CHLOROPHYLL

Fluorescence was recorded continuously using a Turner Model 111 fluorometer as described by Lorenzen (1966). Discrete samples, taken every 30 minutes, were used to calibrate and convert fluorescence to chlorophyll a concentration. The discrete chlorophyll samples were analyzed using the method of Strickland and Parsons (1968). These samples were filtered through a Whatman 4.5 cm G/FC glass fiber filter. For comparison with ATP, chlorophyll concentrations were converted to carbon units (mg/l) by using the average conversion factor (100) proposed by Holm-Hansen (1969).

E. ATP

Adenosine triphosphate was sampled every 10 minutes or approximately every 3 km over most of the cruise tracks. Each 50 ml sample was filtered through a 200- μ m nylon screen, then through a 0.45- μ m glass filter. The method of Holm-Hansen and Karl (1976) was used for ATP analysis. For comparison with chlorophyll, ATP concentrations were converted to carbon units (mg/l) by using the average conversion factor (250) proposed by Holm-Hansen (1969).

F. TEMPERATURE

Sea surface temperature was recorded continuously from a thermistor located at approximately 2.5 m depth coincident to the sea water intake. Sea surface temperature was also measured by bucket thermometer every 20 minutes simultaneously

with the release of a Sippican expendable bathythermograph probe (XBT). The bucket thermometer readings differed by a maximum of $\pm 0.5^{\circ}\text{C}$ with respect to the continuous samples and by a maximum of $\pm 0.6^{\circ}\text{C}$ with respect to the XBT temperatures. When these deviations exceeded $\pm 0.3^{\circ}\text{C}$ they were adjusted to the bucket thermometer readings.

In Figures 8 and 21 the XBT data are plotted as isotherms in vertical sections along the cruise tracks for the 30 April and 7 August cruises. A contour interval of 0.5°C was used.

G. WIND

Wind effects were calculated from the Fleet Numerical Oceanography Center's (FNOC) Data Base using the so-called Field By Information Blending (FIB) routine for grid position $36^{\circ}\text{N} \times 122^{\circ}\text{W}$, which is in the vicinity of the Point Sur coastal region. Computer programs developed by Mr. Andrew Bakun of the National Marine Fisheries Service (Bakun, 1973) produced an output consisting of six hour values for the wind vector, wind stress magnitude, Ekman transport vector, upwelling index (the Ekman vector component normal to the coastline) and vertical velocity (Appendix B). A minimum of 12 days of data prior to and after each cruise was selected for analysis.

IV. RESULTS

A. 30 APRIL CRUISE

Satellite IR imagery indicated a cyclonic thermal feature had developed off Pt. Sur, California, on 19 April 1979 (Plate 1). It appeared to be forming as a seaward extension of coastally upwelled water. By 29 April it appeared to have persisted as a plume-like feature with a cyclonic swirl (Plate 2). Through the use of temperature and nutrient sampling, referenced to the satellite imagery, the central area was found on the first transect and re-located on each subsequent transect (Fig. 1).

Temperature, nitrate and phosphate are plotted against elapsed distance from $36^{\circ}39.1'N$ x $121^{\circ}58.6'W$ (Fig. 2). The feature is evident in the gradients of nitrate, phosphate and temperature.

Correlation coefficients of $r = 0.98$ for nitrate to phosphate, $r = -0.94$ for phosphate to temperature, and $r = -0.96$ for nitrate to temperature were obtained (Table I). Nutrient ratio (N/P) and temperature when plotted against elapsed distance (Fig. 4) also exhibit a strong negative correlation with respect to each other; as is to be expected.

Based on the temperature minima and nutrient maxima the central area of the feature appeared at approximately 65 km (Leg 1), 220 km (Leg 2) and 300 km (Leg 3). The other cold water areas at approximately 165 km (Leg 2) and beyond 335 km




Plate 1. TIROS-N Satellite IR Image of the California Coast, 19 April 1979.




Plate 2. TIROS-N Satellite IR Image of the California Coast, 29 April 1979.




Fig. 1. Track of the 30 April Cruise (solid line) and of the Upwelling Feature (dashed line) based on Satellite IR Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 2. Nitrate, Phosphate, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise. Note the tendency for inverse correlation between temperature and nutrients.




Fig. 3. ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise.




Fig. 4. Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise. Note the tendency for inverse correlation between temperature and nutrient ratio.




Fig. 5. Nitrate Versus Phosphate for the 30 April Cruise.




Fig. 6. Nitrate Versus Temperature for the 30 April Cruise.




Fig. 7. Phosphate Versus Temperature for the 30 April Cruise.




Fig. 8. Vertical Temperature Sections Along the Track of the 30 April Cruise.

(Leg 3) occurred when the ship's track extended into the near-shore upwelling region. Sharp chemical and thermal gradients or "fronts" are evident on all legs.

The linear regression analysis of nitrate versus phosphate, nitrate versus temperature and phosphate versus temperature (Figs. 5, 6, 7) yielded slopes of 15.10, -6.48 and -0.43 respectively and x-axis intercepts of 0.48 μM , 12.99°C and 14.11°C respectively (Table I).

Figure 8 gives some idea of the vertical structure of the upwelled feature. The thermal feature is evident from elapsed distance 64.0 through 81.6 km, 206.2 through 213 km and 312.4 through 323.4 km (where the 11.0°C isotherm surfaces). Coastal upwelling is evident from elapsed distance 163.7 through 171.4 km and 329.3 through 353.8 km.

B. 13 JUNE CRUISE

On 13 June 1979, Mr. Laurence Breaker (NESS, Redwood City) gave approximate coordinates and adjacent shore features of a coastal upwelling event off Point Sur, with "plume-like" characteristics (Plate 3). The ACANIA sailed south parallel to the shore until the northern and southern thermal and nutrient gradients were transected. North-south legs were to be repeated westward in a ladder-like fashion until the seaward termination of the plume was found. However, the cruise was terminated (after only ten hours) due to a clogged sampling port.

Sea surface temperature, nutrient, ATP and chlorophyll data were collected along one and one-half transects of the feature as shown in Fig. 9. The elapsed distances are from point 36°36.3'N x 121°58.7'W.

Based on the distributions of temperature, nitrate and phosphate (Fig. 10) the upwelling area was encountered at approximately 15 km along the track. Nitrate and phosphate values were high (up to 12.1 μ M and 1.23 μ M respectively) until the southern frontal boundary was reached. In the oceanic water, at approximately 50 km, nitrate and phosphate concentrations were low (0.55 to 3.8 μ M and 0.33 to 0.65 μ M, respectively). To the north, the frontal southern boundary was again encountered at approximately 60 km along the track. Throughout, there was a strong correlation ($r = 0.93$) between nitrate and phosphate, a strong inverse correlation between nitrate and temperature ($r = -0.92$) and a strong inverse correlation between temperature and phosphate ($r = -0.93$).

The chlorophyll data are unreliable after an elapsed distance of 40 km, because of excessive air injection into the sea chest during rough seas. Additionally, the XBT recorder was inoperative during this cruise.

C. 7 AUGUST CRUISE

The satellite image of 30 July 1979 (Plate 4) shows a cold water plume extending approximately 150 km southwest from Point Sur. Mr. Laurence Breaker (NESS, Redwood City)




Plate 3. TIROS-N Satellite IR Image of the California Coast, 13 June 1979.




Fig. 9. Track of the 13 June Cruise (solid line) and Outline of the Upwelling Feature (dashed line) Based on Satellite Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 10. Nitrate, Phosphate, Nutrient Ratio, ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 13 June Cruise.




Fig. 11. Nitrate Versus Phosphate for the 13 June Cruise.




Fig. 12. Nitrate Versus Temperature for the 13 June Cruise.




Fig. 13. Phosphate Versus Temperature for the 13 June Cruise.

reported the feature had continued to extend southwest. By 5 August 1979 the feature had extended approximately 240 km southwest of Point Sur (Plate 5). Sea surface temperatures were expected to be within approximately 14 to 15°C inside the plume and greater than 17°C outside the plume based on ship reports correlated to the satellite image by Mr. Breaker.

The cruise plan was similar to the 30 April and 13 June cruises except, in this case, the primary interest was the water near the seaward termination of the feature. The track strategy was planned to coincide with the axis of the feature; when applicable, turn south to transect its southern boundary, zigzag across its southern boundary, then run back up the axis upon return. Due to limited sampling resources, ATP sampling did not commence until the southern boundary was reached.

The cruise track is shown in Fig. 14. Legs 2 through 5 intersected the feature's southern boundary. Elapsed distances are measured from position 36°38.8'N x 121°57.5'W.

Figure 15 is the line graph of temperature, nitrate and phosphate along the cruise track. The nutrient lines are discontinuous from 160 to 200 km, 275-295 km and 325-355 km because the Autoanalyzer was secured for cleaning. When there is a phosphate line but no line for nitrate, nitrate concentrations were so low that they were below the sensitivity of the instrument. However, in all cases the frontal crossings are clearly evident in the temperature lines.

Frontal crossings occurred at approximately 260, 325, 360 and 385 km on Legs 2 through 5. The line graphs for Legs 1 and 6 are generally within a temperature range of 14 to 16°C, indicating that the ACANIA was within the plume.

The correlation coefficients were $r = 0.42$ for nitrate versus phosphate, $r = -0.11$ for nitrate versus temperature, and $r = -0.39$ for phosphate versus temperature. Linear regression analysis yielded slopes of 26.44, -13.63 and -0.32 and x-axis intercepts of 0.61 μM phosphate, 15.35°C and 17.56°C for Figs. 18, 19, and 20, respectively (Table I).

The thermal feature is evident in the vertical cross section of temperature (Fig. 21) and gives some idea of the vertical structure of the feature. The transition between plume water and oceanic water is located between elapsed distances 249.5 through 263.1 km, 274.8 through 285.9 km, and 305.2 through 336.0 km (where the 15.5°C isotherm rises toward the surface). The 15.5°C isotherm is 30 to 40 meters deep outside the plume and inside it rises toward the surface. The thermal patchiness, evident in the satellite IR imagery, is seen in both the temperature cross section and the surface temperature line graphs.




Plate 4. TIROS-N Satellite IR Image of the California Coast, 30 July 1979.




Plate 5. TIROS-N Satellite IR Image of the California Coast, 5 August 1979.




Fig. 14. Track of the 7 August Cruise (solid line) and Outline of the Upwelling Feature (dashed line) based on Satellite Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 15. Nitrate, Phosphate, and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise.




Fig. 16. ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise.




Fig. 17. Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise Track.




Fig. 18. Nitrate Versus Phosphate for the 7 August Cruise.




Fig. 19. Nitrate Versus Temperature for the 7 August Cruise.




Fig. 20. Phosphate Versus Temperature for the 7 August Cruise.




Fig. 21. Vertical Temperature Sections Along the Track of the 7 August Cruise

TABLE I
SUMMARY OF REGRESSION ANALYSES

CRUISE	\bar{N} (μM)	\bar{P} (μM)	\bar{T} ($^{\circ}\text{C}$)	SLOPE			"X" INTERCEPT			CORRELATION COEFFICIENTS		
				$\Delta N:\Delta T$	$\Delta N:\Delta T$	$\Delta N:\Delta P$	$N:T(^{\circ}\text{C})$	$P:T(^{\circ}\text{C})$	$N:P(\mu\text{M})$	$N:T$	$P:T$	$N:P$
April 30 to May 1	8.59	1.03	11.7	-6.48	-0.43	15.1	12.99	14.11	0.48	-0.96	-0.94	0.98
June 13	8.24	0.90	12.4	-4.99	-0.39	12.88	13.79	14.43	0.26	-0.92	-0.93	0.93
August 7 to 9	5.28	0.767	15.06	-13.63	-0.32	26.44	15.35	17.56	0.61	-0.11	-0.39	0.42

TABLE II
DATA RANGES

TEMPERATURE °C	NITRATE μM	PHOSPHATE μM	RATIO N/P	ATP ng/l	CHLOROPHYLL mg/m ³	ATP mgC/l	CHLOROPHYLL mgC/ℓ
<u>30 April to 1 May Cruise</u>							
9.4 to 13.6	0.29 to 23.11	0.45 to 2.06	0.6:1 to 11.7:1	338 to 4126	0.13 to 6.49	0.0845 to 1.0315	0.013 to 0.649
<u>13 June Cruise</u>							
11.1 to 13.8	0.55 to 12.10	0.33 to 1.29	1.7:1 to 12.3:1	825 to 2485	1.15 to 17.28	0.206 to 0.621	0.115 to 1.728*
<u>7 to 9 August Cruise</u>							
12.2 to 17.2	none detected to 20.91	0.03 to 2.07	0.0 to 15.3:1	712 to 3418	0.17 to 8.19	0.178 to 0.855	.017 to 0.819

*Data are unreliable due to air injection into the sampling water.

V. DISCUSSION

A better understanding of the "life cycle" of an upwelling event may be provided through the integration of satellite IR imagery and in situ sampling. This approach leads to a descriptive "phase" model of the formation and dissipation of an upwelling chemical system and some insight on the interaction of dynamic and biological processes in the system. This thesis analyzes results obtained while using this integrated technique to investigate three upwelling events.

An upwelling event, like many physical phenomena, may be described in terms of phases: initiation, growth, equilibrium, decay and dissipation. Each phase is a function of the magnitude and rate of change of the dynamic processes (mixing, advection, diffusion) and the boundary conditions (atmospheric forcing; topography and bathymetry). The chemical characteristics of any phase of an upwelling event are a function of the initial conditions (source water); the magnitude, direction, and duration of the driving function (wind stress); and the differential utilization and regeneration by biological processes.

The satellite imagery prior to all cruises show "plumes" of colder water extending from the coast near Point Sur. These features range in scale from tens of kilometers to several hundred kilometers in length. Analysis of in situ

surface nutrient concentrations, summarized in Tables I and II, the surface winds in the area prior to and during the cruises (Appendix B), sea surface temperatures, satellite IR imagery and bathymetry strongly argue that these satellite detected thermal plumes were caused by upwelling events. That nutrient concentrations may closely follow the pattern of the satellite thermal imagery of the sea surface was shown by Traganza et al. (1980). These upwellings of cold nutrient-rich waters produce chemical and thermal fronts which appear to have associated with them increased biological activity.

The feature examined during the 30 April Cruise is postulated to be composed of recently upwelled water. Figure 2 shows very sharp chemical (up to $\frac{10.41 \mu\text{M NO}_3}{4.7 \text{ km}}$) and thermal (up to $0.9^\circ\text{C}/2.3 \text{ km}$) gradients. The range of nutrient ratio values are 0.6N:1P (lowest value, outside the feature) and 11.7N:1P (maximum level, inside the feature). The graph of nutrient ratio (Fig. 4) against elapsed distance along the track shows that the concentrations of high values are either within the area of the feature or the coastal upwelling. The low values of temperature (below 12°C) and the high concentrations of nitrate and phosphate (up to $12.1 \mu\text{M}$ and $1.29 \mu\text{M}$, respectively) are all either within the feature or the coastal upwelling (Fig. 10). There are strong correlations between nitrate and phosphate ($r = 0.98$), nitrate and temperature ($r = 0.96$), and phosphate and temperature ($r = -0.94$). Analysis of the wind data prior to the

cruise shows that favorable upwelling conditions occurred during the periods 9 to 20 April and 28 April to 4 May (Appendix B). All these factors, plus the lower temperatures indicated by the satellite IR imagery are taken to describe a feature of recently upwelled water in the early phases of its life cycle.

The 13 June Cruise examined a feature that is close inshore (Plate 3). Wind data (Appendix B) shows favorable upwelling conditions commenced only three days prior to cruise. The waters outside the feature are about 14°C but inside the feature temperatures fall below 12°C. All high concentrations of nutrients are within the feature, as are all high nutrient ratios. There are strong correlations between nitrate and phosphate ($r = 0.93$), nitrate and temperature ($r = -0.92$), and phosphate and temperature ($r = -0.93$) (Figs. 11, 12 and 13). This feature is recently upwelled water and is postulated to be in the early phases of the upwelling event "life cycle."

The waters examined in these two cruises still have very strong source water (thermal and nutrient) signatures. The dynamic and biological processes have not had sufficient time to change the upwelled water's initial characteristics. Along the frontal boundaries, the biological processes become evident (Figs. 3 and 10). The dynamic processes appear to have provided the necessary environment for biological activity.

The data from the 7 August Cruise were gathered predominantly from within the feature (Fig. 14). In comparison to the previous cruises, this resulted in a considerably smaller proportion of data points being collected outside the feature. A statistically insufficient number of samples were obtained in the low nutrient concentration, high temperature region. In comparison to the previous cruises the analysis is biased as follows: mean nutrient concentrations are high; mean temperature is low; the slopes of nitrate to phosphate, nitrate to temperature, and phosphate to temperature are too high, and the respective x-axis intercepts are too low.

The feature observed on the 7 August Cruise appears to contain upwelled water that is in various phases of the upwelling cycle. At 50 km elapsed distance water temperature was below 13°C with rapidly increasing nutrient ratios. Sampling was interrupted to clean the Autoanalyzer but concentrations were approaching levels consistent with the previous cruises (nutrient ratios above 10N:1P). This area of the feature contains water that has source water characteristics and may be assumed to be in the early phases. At 75 km elapsed distance, while still within the feature, temperature was approximately 14°C and nutrients had fallen, but not to the "outside-the-features-levels" of the previous cruises. This area might be considered to be close to equilibrium. The water was still relatively cold and nutrient-rich but no longer similar to the source water; the

dynamic and biological processes apparently had started to modify it. The temperature then rose to the 15.0° to 15.5°C range and while nutrients still generally decreased with increasing temperature, they showed considerable variability. This area can be characterized as in the decay phase. The dynamic and biological processes continued to change the temperature and nutrient characteristics.

Once outside the feature, temperatures rose sharply to levels of 17.5°C and nutrients correspondingly fell. As the track moved west (Fig. 15) the nutrient levels and nutrient ratios (Fig. 17) never recovered to the levels previously encountered within the feature. In some areas, at the western most extent of the cruise, the nitrate levels are too low to be detected. In most coastal upwelling systems, nitrogen is characteristically the limiting nutrient, Dugdale et al. (1967) and Thomas, (1969). "In accordance with Liebig's law of the minimum, that constituent of the sea water present in smallest quantity relative to the requirement for growth of organisms will become the limiting factor." (Redfield, 1958). In this area, there is insufficient nitrate to sustain photosynthesis and, therefore, the biological modification of the feature is complete and low biomass concentrations are evident (Fig. 16). All that remains is for the dynamic processes to dissipate the temperature anomaly.

The wind data prior to the cruise shows a cyclical pattern of upwelling events occurring since early April.

Upwelling periods were of the order of one to three weeks and relaxation periods were of the order of two to six days (Appendix B). The average Ekman transport varied from west southwest to southwest. With such an intermittent, long range pattern of offshore transport; it is therefore possible that the feature investigated on the 7 August Cruise was in fact upwelled water of varying ages. Barton, et al. (1977) indicates that during an upwelling event coldest water will migrate to the continental shelf edge and remain there until the system relaxes. Upon return of favorable winds the upwelling will again reappear along the inner shelf. In this case, high nutrient, cold water was found at 50 km elapsed distance which is just inside the shelf break. The center of an upwelling may stop at the shelf edge, with continued favorable winds, upwelled water will continue to be transported offshore in a 20 m thick surface layer (Huyer, 1974). Patchiness or banding can result from this start/stop action between close upwelling events (Barton, et al. 1977). The satellite images (Plates 4 and 5) show some patchiness. The second image (Plate 5) shows the feature has moved 90 km further southwest and the wind data show the Ekman transport to be west southwesterly during the period between the images. Therefore the feature investigated has some of the characteristics of a wind induced upwelling plume with the oldest water at its furthest extent; still maintaining some of its temperature characteristics but due to biological action and mixing losing much of its high nutrient signature at its westernmost (oldest) edges.

In viewing the correlation values of all cruises it must be remembered "that where elements are substantially depleted by the growth of plants small unused residues of one or another element may greatly alter the ratios." (Redfield, 1958). The nutrient-to-temperature correlations will reflect the effects of heat transfer, mixing, advection, diffusion and biological utilization and regeneration. The longer the water has been on the surface, the more time the processes have to affect the correlations. This is evident on the 7 August Cruise where all correlations are poor.

In sea water the ratio of change of nitrate and phosphate, $\Delta N/\Delta P$, is representative of the uptake of these nutrients by phytoplankton at a rate of 16N:1P. The slope of the best fit line found from linear regression analysis (Table 1) represents this ratio of change. The values obtained on the 30 April Cruise (15.1:1) and 13 June Cruise (12.88:1) are close to the theoretical value of 16N:1P.

On all cruises the nutrient ratio did not equal the theoretical value of 16:1 (Redfield, 1958) most likely because the source water was not at that ratio. The ratios ranged from a low where no nitrate was detected to a high of 15.3:1. The low values perhaps were found where further biological uptake had become inhibited due to low nitrate concentration and the higher values corresponded to the upwelled water. These values are not inconsistent with the annual range for the nutrient ratio of 3:1 to 13:1 obtained in a study by Butler et al. (1979).

Finally, in areas of upwelling there exist some geographically fixed preferred positions, e.g. south of capes, where plumes of freshly upwelled water protrude offshore (Shaffere, 1976; Reid et al., 1958). The research of this thesis and that of Traganza et al. (1980), which covered one annual cycle, indicates the coastal waters off Point Sur, California, are such an area. Whether this is because of the proximity of Point Sur and/or Monterey Canyon, local bathymetry, local water circulation patterns, local wind effects, or coastally trapped topographic Rossby waves; or a combination of these factors cannot be determined from the data. However, offshore plumes consistently, albeit aperiodically, originate from this area.

VI. CONCLUSIONS

1. High resolution, enhanced satellite IR imagery can be of great value in locating and assessing feature motion when investigating upwelling events that are manifested as sea surface temperature anomalies.
2. Upwelled water may be evidenced at considerable (ca. 100 to 300 km) distances from the coast.
3. Qualitative inferences on the distribution of nutrient concentrations can be made using satellite infrared imagery if coupled with in situ sampling to establish nutrient versus temperature correlations.
4. Altogether, with sufficient historical upwelling data from in situ monitoring, a nutrient predictive model for upwelled water in the initiation, growth and equilibrium phases is conceivable in the future.
5. A predictive model of the entire upwelling cycle, in all its phases, is not yet possible. The effects of air-sea interaction, the dynamical processes, and the biological processes make any nutrient model extremely complex. The dynamic processes vary from region to region and season to season. Regeneration and utilization rates vary from organism to organism and season to season. This further compounds the task.
6. The area off Point Sur, California, consistently has offshore upwelling plumes and merits further detailed

study. Such study is continuing under the direction of Dr. Eugene Traganza supported by the Office of Naval Research, Code 482.

7. From a Naval aspect the thermal gradients in upwelled water can have a considerable impact on sound propagation. The increased biological concentrations associated with the thermochemical fronts can, depending on the species, increase the reverberation levels and background noise.

64

TIME GMT 1604	LATITUDE NORTH 36 39.1	LONGITUDE WEST 121 58.6	DISTANCE KM 0.0	NO3 UM 0.48	PO4 UM 0.53	NUTR.RATIO NO3/PO4 0.9	ATP NG/L 2648.0	CHL A MG/100 5.57	TEMP DEG C 12.0
			4.7	0.83	0.50	1.5		5.06	12.6
			5.3	0.65	0.54	1.2		4.94	12.4
			6.4	1.67	0.76	4.8		4.34	11.9
			7.0	1.36	0.61	2.2		4.35	12.2
			8.2	0.61	0.55	2.2		4.18	12.3
			8.3	0.83		1.5		4.29	12.4
			9.3					4.42	12.4
			10.5					4.65	12.4
			11.7	1.54	0.56	2.8		4.31	12.4
			12.3	2.60	0.65	4.0		4.99	12.4
			13.4	5.62	0.86	6.5		4.56	12.0
			14.0	7.21	1.01	7.5		4.03	11.6
			15.2	7.75	1.22	8.8		3.53	11.5
			16.3	10.41	1.19	8.8		2.59	11.3
			17.5	9.34	1.12	7.8		2.49	11.2
			18.7	10.76	1.19	8.5		1.59	11.3
			19.8	9.16	1.21	8.5		1.30	11.3
			21.0	10.23	1.22	8.4		1.01	11.3
			22.2	10.58	1.22	8.7		1.02	11.4
			23.3	10.23	1.22	8.5		1.37	11.5
			24.5	10.05	1.19	8.4		1.41	11.6
			25.7					1.17	11.7
			26.8					1.11	11.8
			27.4					1.14	11.9
			28.6					1.12	12.0
			29.8					1.11	12.1
			30.9					1.12	12.1
			31.5					1.12	12.1
			32.7					1.07	12.1

R/V ACANIA

30 APR / 1 MAY 79

CHEMICAL MEASUREMENTS (CRUISE VI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	MCS UM	PM4 U/L	NUTR. RATIO PM4/PM4	ATP NG/L	CHL. A MG/L	TEMP DEG C
2000	36 21.6	122 6.9	33.3	6.68	0.95	7.0	2078.0	3.67	12.1
			33.3	6.86	0.97	7.1		2.67	12.0
			34.4	7.21	0.98	7.3		2.86	12.0
			35.0	7.04	0.97	7.0		2.92	12.0
			35.6	6.86	1.00	7.0		1.58	12.0
			36.2	7.04	0.98	6.6	1823.0	1.21	12.0
			36.3	6.50	0.98	6.5		0.82	11.9
			37.3	6.15	0.97	6.0		0.84	12.1
			37.9	5.62	0.93	5.0		0.86	12.1
			38.5	4.55	0.87	5.2		0.87	12.2
			39.1	4.30	0.84	5.0	1192.0	0.88	12.4
			39.7	3.85	0.81	4.8		0.90	12.4
			40.3	4.02	0.79	5.1			12.4
			40.8	3.67	0.78	4.7			12.5
			41.4	3.14	0.75	4.2			12.5
			42.0	2.96	0.73	4.1	1146.0		12.6
			42.6	2.96	0.73	4.1			12.6
			43.2	2.96	0.73	4.1			12.7
			43.8	2.96	0.73	4.1			12.7
			44.3	2.96	0.73	4.1			12.7
			44.9	2.78	0.68	4.1			12.7
			45.5	3.14	0.72	4.4	1186.0		12.7
			46.1	3.67	0.76	4.8			12.7
			46.7	4.33	0.81	5.4			12.7
			47.3	5.44	0.89	6.2			12.7
			47.9	7.21	1.00	7.5	1508.0	2.51	12.7
			48.4	8.10	1.03	7.8			12.7
			49.0	8.90	1.11	8.0			12.7
			49.6	8.99	1.11	8.1			12.7
			50.2	9.52	1.15	8.3			12.7
			50.8				1399.0	0.90	12.7
			51.4				1286.0		12.7
			52.0						12.7
			52.6	11.86	1.30	9.1			12.7
			53.2	12.07	1.33	9.1			12.7
			53.8	13.05	1.36	9.3	1246.0		12.7
			54.4	13.88	1.53	10.7			12.7
			55.0	16.99	1.56	10.6			12.7
			55.6	17.77	1.61	10.8	1578.0	1.46	12.7
			56.2	18.01	1.66	10.9		1.33	12.7
			56.8	18.25	1.67	11.0		1.07	12.7
			57.4	18.31	1.69	11.0	1413.0	0.94	12.7
			58.0	18.80	1.70	11.1		0.81	12.7
			58.6	19.23	1.73	11.1			12.7
			59.2	19.46	1.77	11.2			12.7
			59.8	20.11	1.78	11.3			12.7
			60.4	20.98	1.78	11.3			12.7
			61.0	17.01	1.56	10.9	1796.0	1.11	12.7
			61.6						12.7
			62.2						12.7
			62.8						12.7
			63.4						12.7
			64.0						12.7
			64.6						12.7
			65.2						12.7
			65.8						12.7
			66.4						12.7
			67.0						12.7
			67.6						12.7
			68.2						12.7
			68.8						12.7
			69.4						12.7
			70.0						12.7
			70.6						12.7
			71.2						12.7
			71.8						12.7
			72.4						12.7
			73.0						12.7
			73.6						12.7
			74.2						12.7
			74.8						12.7
			75.4						12.7
			76.0						12.7
			76.6						12.7
			77.2						12.7
			77.8						12.7
			78.4						12.7
			79.0						12.7
			79.6						12.7
			80.2						12.7
			80.8						12.7
			81.4						12.7
			82.0						12.7
			82.6						12.7
			83.2						12.7
			83.8						12.7
			84.4						12.7
			85.0						12.7
			85.6						12.7
			86.2						12.7
			86.8						12.7
			87.4						12.7
			88.0						12.7
			88.6						12.7
			89.2						12.7
			89.8						12.7
			90.4						12.7
			91.0						12.7
			91.6						12.7
			92.2						12.7
			92.8						12.7
			93.4						12.7
			94.0						12.7
			94.6						12.7
			95.2						12.7
			95.8						12.7
			96.4						12.7
			97.0						12.7
			97.6						12.7
			98.2						12.7
			98.8						12.7
			99.4						12.7
			100.0						12.7

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE K	NCS UM	PN4 UM	NUTR. RATIO NO3/PN4	ATP MG/L	CHL A MG/M3	TEMP DEG C
2200	36	3.7	122	0.2					
			67.1	14.97	1.44	10.4		4.71	10.2
			67.1	13.05	1.37	10.3		4.13	10.4
			68.3	13.71	1.37	10.0		3.89	10.8
			69.0	13.14	1.34	10.3		4.10	10.0
			70.0	13.21	1.30	10.2		4.29	11.1
			70.6	13.04	1.22	9.8		5.88	11.3
			71.2	11.92	1.22	9.8		4.90	11.4
			71.8	11.79	1.23	9.6		4.13	11.4
			72.3	11.82	1.22	9.7		4.10	11.4
			73.5	11.85	1.22	9.3		3.89	11.3
			74.1	11.11	1.20	9.8		4.29	11.3
			74.7	11.12	1.22	9.8		5.03	11.6
			75.3	11.56	1.19	9.7		6.54	11.5
			75.8	11.01	1.15	9.6		5.88	11.6
			76.4	11.03	1.15	9.6		6.54	11.5
			77.0	7.22	0.95	7.6		6.54	11.5
			77.6	7.04	0.97	8.1		5.89	11.6
			78.2	6.86	0.86	7.0		5.89	11.5
			79.3	6.05	0.78	5.9		5.89	11.5
			80.5	3.59	0.73	4.9		5.89	11.5
			81.1	3.18	0.70	4.5		5.89	11.5
			81.7	2.84	0.65	4.4		5.89	11.5
			82.3	2.27	0.62	4.4		5.89	11.5
			83.0	2.97	0.68	4.7		5.89	11.5
			84.0	3.18	0.70	4.9		5.89	11.5
			85.0	3.19	0.70	4.9		5.89	11.5
			86.4						
			87.5						
			88.7						
			89.3						
			90.4						
			91.0						
			91.6						
			92.8						
			93.9						
			94.5						
			95.7						
2300	35	56.2	121	53.5					
			86.4	2.12	0.48	4.4		4.71	12.0
			87.5	2.54	0.64	4.0		4.13	12.0
			88.7	2.97	0.67	4.4		4.29	12.0
			89.3	2.76	0.65	4.4		4.10	12.0
			90.4	2.11	0.61	4.4		4.29	12.0
			91.0	2.10	0.61	4.4		4.29	12.0
			91.6	2.54	0.64	4.0		4.13	12.0
			92.8	2.32	0.61	4.4		4.29	12.0
			93.9						
			94.5						
			95.7						

R/V ACANIA

30 APR / 1 MAY 79

CHEMICAL MESOSCALE(CRUISE VI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	PH3 M	PO4 M	NUTR RATIO PO3/PO4	ATP NG/L	CHL A MG/73	TEMP DEG C
			125.4	1.01	0.59	1.7		1.15	13.0
			126.0	1.19	0.60	2.0		1.29	13.0
			127.0	1.19	0.60	2.0		1.41	13.0
			127.7	1.19	0.62	1.9	1023.0	1.54	12.9
			128.4	1.01	0.61	1.7		1.34	12.9
			128.9	1.01	0.59	1.7		1.49	12.9
			129.5	1.01	0.62	1.6	1773.0	1.71	12.8
			130.1	0.83	0.53	1.6		1.77	12.8
			130.7	0.47	0.53	0.9		1.58	12.5
			131.3	0.65	0.53	1.2		1.56	12.9
			131.9	0.47	0.50	0.9		1.46	12.9
			132.4	0.29	0.49	0.6	1121.0	1.23	13.2
			133.0	0.0	0.49	0.0		1.11	13.2
			133.6	0.0	0.49	0.0		0.99	13.2
			134.2	0.0	0.49	0.0		0.87	13.2
			134.8	0.0	0.51	0.0	1264.0	0.75	13.3
			135.4	0.0	0.51	0.0		0.85	13.3
			136.5	0.0	0.51	0.0		0.86	13.3
			137.1	0.0	0.51	0.0		0.83	13.2
			137.7					0.92	13.2
			138.3				1051.0	0.90	13.2
			139.9					0.88	13.2
			140.0					0.87	13.2
			140.6					0.85	13.2
			141.2				1297.0	0.84	13.2
			141.8					0.85	13.2
			142.4	0.47	0.58	0.8		0.85	13.2
			142.9	0.29	0.53	0.5		0.85	13.2
			143.5	0.47	0.53	0.9		0.86	13.2
			144.1	0.47	0.53	0.9	1397.0	0.97	13.2
			144.7	0.65	0.52	1.3			13.2
			145.3	0.47	0.52	0.9			13.2
			146.4	0.29	0.52	0.6			13.2
			147.0	0.29	0.52	1.2		1.56	13.1
			147.6	0.65	0.54	1.2		1.40	13.1
			148.2	0.65	0.56	1.2		1.31	13.1
			148.8	0.47	0.57	0.8		1.25	13.0
			149.4	0.47	0.64	0.8	2895.0	1.17	12.9
			149.9	1.55	0.72	0.9		1.18	12.9
			150.5	2.09	0.72	0.9		1.16	12.9
			151.1	2.27	0.72	0.9		1.12	12.8
			152.3	2.63	0.75	0.9		1.19	12.6
			152.9	3.17	0.83	0.8		0.94	12.5
			153.4	3.35	0.83	0.8		1.07	12.4
			154.0		0.82	0.8	1444.0		

0200 35 36.8 121 31.7

[illegible]

R/V ACANIA										30 APR / 1 MAY 79										CHEMICAL MESUSCALE(CRUISE VI)									
TIME GMT	LATITUDE NORTH		LONGITUDE WEST		DISTANCE KM		NT3 HM		PO4 UM		NUTP-RATIO NT3/PO4		ATP NG/L		CHL A MG/K3		TFWD NEG C												
0500	35	57.7	121	41.8	183.8	5.88	0.85	6.9																					
					184.4	5.83	0.87	6.8																					
					185.0	5.34	0.82	6.8																					
					186.1	4.62	0.79	5.8																					
					186.7	4.26	0.76	5.8																					
					187.3	4.08	0.74	5.5																					
					187.9	3.72	0.71	5.2																					
					188.5	3.54	0.71	5.0																					
					189.0	3.72	0.73	5.1																					
					190.2	4.08	0.75	5.4																					
					191.4	4.90	0.78	6.2																					
					192.0	4.76	0.77	5.7																					
					192.5	3.90	0.74	5.3																					
					193.1	3.72	0.72	5.2																					
					193.7	3.54	0.73	5.4																					
					194.3	4.08	0.75	5.7																					
0600	36	1.7	121	51.8	194.3	4.08	0.81	5.9																					
					195.5	4.80	0.81	5.9																					
					196.0	4.80	0.81	5.9																					
					196.6	4.44	0.78	5.7																					
					197.2	4.26	0.76	5.6																					
					197.8	4.44	0.80	5.5																					
					198.4	4.62	0.81	5.7																					
					199.0	4.90	0.78	5.9																					
					199.5	4.44	0.78	5.7																					
					200.1	4.44	0.78	5.7																					
					200.7	2.99	0.72	4.2																					
					201.3	2.81	0.72	3.9																					
					201.9	2.45	0.69	3.6																					
					202.5	3.17	0.74	4.3																					
					203.0	3.45	0.77	4.3																					
					203.6	3.90	0.78	5.0																					
204.2	3.90	0.78	5.0																										
204.8	3.90	0.78	5.0																										
205.4	3.54	0.75	4.7																										
206.0	3.17	0.73	4.3																										
206.5	3.35	0.73	4.3																										
207.1	3.54	0.80	4.9																										
207.7	3.54	0.77	4.6																										
208.3	5.34	0.88	6.1																										
208.9	6.78	0.99	6.8																										
209.5	7.68	1.05	7.3																										
210.0																													
210.6																													
211.2																													
211.8																													
212.4																													
														2066.0															

P/V ACANIA									
30 APR / 1 MAY 79									
CHEMICAL MESOSCALE(CRUISE VI)									
TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PD4 UM	NUTR-RATIO NO3/PD4	ATP NG/L	CHL A MG/M3	TEMP DEG C
0700	36	5.2	122	1.6	213.0	1.30	10.2	1.96	11.1
					213.5	1.36	10.3	2.23	11.1
					214.1	1.39	10.6	2.17	10.9
					214.7	1.44	10.6	2.61	10.6
					215.3	1.45	10.8	1.97	10.5
					215.9	1.54	11.0	1.54	10.4
					216.5	1.60	11.0	1.30	10.2
					217.0	1.65	11.2	0.94	10.0
					217.6	1.68	11.2	0.75	9.9
					218.2	1.73	11.2	0.74	9.9
					219.8	1.71	11.2	0.74	9.9
					219.4	1.77	11.2	0.74	9.9
					220.0	1.76	11.0	0.74	9.9
					220.5	1.76	11.0	0.74	9.9
					221.1	1.73	11.1	0.96	10.0
					221.7	1.72	11.1	0.96	10.0
					222.3	1.70	11.1	0.99	10.0
					222.9	1.71	10.9	0.90	10.0
					223.5	1.69	11.0	0.71	10.0
					224.0	1.68	10.8	0.15	10.0
					224.6	1.66	10.7	0.89	10.0
					225.2	1.66	10.7	0.15	10.0
					225.8	1.70	10.5	0.83	10.0
					226.4	1.70	10.5	0.98	10.0
					227.0	1.63	10.4	0.81	10.0
					227.5	1.57	10.5	0.94	10.0
					228.1	1.54	10.4	0.81	10.0
					228.7	1.53	10.3	0.81	10.0
					229.3	1.48	10.4	1.41	10.0
					229.9	1.47	10.3	1.47	10.0
					230.5	1.46	10.2	1.47	10.0
					231.0	1.45	10.2	1.53	10.0
					231.6	1.44	10.2	1.53	10.0
					232.2	1.43	10.2	1.53	10.0
					232.8	1.42	10.2	1.53	10.0
					233.4	1.41	10.2	1.53	10.0
					234.0	1.40	10.2	1.53	10.0
					234.6	1.39	10.2	1.53	10.0
					235.1	1.38	10.2	1.53	10.0
					235.7	1.37	10.2	1.53	10.0
					236.3	1.36	10.2	1.53	10.0
					236.9	1.35	10.2	1.53	10.0
					237.5	1.34	10.2	1.53	10.0
					238.0	1.33	10.2	1.53	10.0
					238.6	1.32	10.2	1.53	10.0
					239.2	1.31	10.2	1.53	10.0
					239.8	1.30	10.2	1.53	10.0
					240.4	1.29	10.2	1.53	10.0
					241.0	1.28	10.2	1.53	10.0
					241.5	1.27	10.2	1.53	10.0

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NU3 UM	PO4 UM	NU3/P04 RATIO	ATP NG/L	CHL A MG/M3	TEMP DEG C
0800	36	9.1	122 12.6	242.1	1.17	0.67	684.0	0.65	12.0
				242.7	2.32	0.67		0.65	12.0
				243.3	2.35	0.59		1.18	12.0
				243.9	1.87	0.55		0.41	12.4
				244.5	1.50	0.54		1.69	12.3
				245.0	1.12	0.52		0.46	12.4
				245.6	1.11	0.50		0.75	12.4
				246.2	0.91	0.50		0.64	12.0
				246.8	0.90	0.50		0.53	12.0
				247.4	0.07	0.50		0.32	12.0
				248.0	1.03	0.56	564.0	0.53	12.0
				248.5	2.18	0.57		0.59	12.0
				249.1	2.37	0.60		0.48	12.0
				250.3	2.55	0.60		0.52	12.0
				250.9	2.54	0.62		0.37	12.0
				251.5	2.15	0.60		0.41	12.0
				252.1	1.55	0.58		0.64	12.0
				253.2	1.53	0.52		0.45	12.0
				253.8	1.71	0.55		0.45	12.0
				254.4	1.10	0.58		0.23	13.1
0900	36	12.2	122 23.2	255.0	2.68	0.59	338.0	0.22	13.2
				255.6	2.68	0.59		0.22	13.2
				256.1	2.67	0.59		0.22	13.2
				257.3	2.66	0.59		0.22	13.3
				257.9	2.66	0.58		0.22	13.3
				258.5	2.45	0.58		0.21	13.0
				259.1	2.45	0.58		0.21	13.0
				260.8	2.23	0.58		0.21	13.4
				261.4	2.02	0.55		0.37	13.4
				262.0	2.01	0.56		0.37	13.4
				262.6	2.01	0.56	446.0	0.54	13.0
				263.1	1.79	0.56		0.24	13.0
				263.7	1.58	0.54		0.24	13.0
				264.3	1.57	0.54		0.20	13.0
				264.9	1.36	0.54		0.20	13.0
				265.5	1.35	0.52		0.20	13.1
				266.1	1.34	0.53		0.20	13.1
				266.6	0.91	0.53		0.20	13.3
				267.2	0.90	0.48		0.13	13.5
				267.8	0.68	0.47		0.15	13.5
1100	35	52.9	122 14.3	268.4	0.47	0.47	342.0	0.14	13.6
				269.0	0.45	0.45		0.14	13.6
				269.6	0.44	0.45		0.20	13.4
				270.1	0.42	0.45		0.20	13.4
				270.9	0.38	0.45		0.32	13.4
				271.3	0.38	0.45		0.32	13.4
				271.9	0.38	0.45		0.32	13.4
				272.5	0.38	0.45		0.32	13.4
				273.1	0.38	0.45		0.32	13.4
				273.7	0.38	0.45		0.32	13.4

R/V ACANIA

30 APR / 1 MAY 79

CHEMICAL MESOSCALE(CRUISE VI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NDJ UM	PD4 UM	NOTE: RATIO NDJ/PO4	ATP MG/L	CHL A MG/3	TEMP DEG C
1200	35 59.6	122 6.0	314.5	10.13	1.10	9.2		2.06	11.5
			315.1	10.50	1.13	9.3		2.06	11.6
			315.6	10.31	1.10	9.4		2.06	11.5
			316.2	10.13	1.10	9.3		2.06	11.3
			317.4	9.94	1.11	9.0		1.90	11.4
			318.0	6.42	0.87	7.4		1.87	11.4
			318.6	5.12	0.84	6.1		1.87	11.4
			319.1	8.69	0.99	8.2		1.90	11.4
			319.7	10.31	1.14	9.0	1701.0	2.27	11.4
			320.3	12.91	1.26	10.2		2.27	11.7
			320.9	12.35	1.25	9.9		2.27	11.5
			321.5	12.72	1.23	9.9		2.27	10.9
			322.1	13.65	1.33	10.3	1155.0	2.23	10.9
			322.6	14.39	1.30	11.1		2.23	10.9
			323.0	14.21	1.33	10.3		2.23	10.7
			324.4	14.39	1.33	10.3		2.23	10.7
			325.0	15.69	1.47	10.7		2.23	10.6
			325.6	15.03	1.66	11.5	1119.0	2.24	10.6
			326.1	15.58	1.71	11.4		2.24	10.5
			326.7	19.77	1.73	11.4		2.24	10.5
			327.3	20.14	1.76	11.4		2.24	10.5
			327.9	17.92	1.60	11.2	948.0	2.23	10.1
			328.5	17.36	1.56	11.1		2.23	10.1
			329.1	16.25	1.51	10.8		2.23	10.3
			329.6	14.53	1.42	10.3		2.23	10.4
			330.2	13.84	1.36	10.2	2055.0	2.23	10.5
			330.9	11.72	1.20	9.4		2.20	10.5
			332.0	11.24	1.13	8.6		2.20	10.6
			332.6	9.76	1.13	9.1	3179.0	2.22	10.8
			333.2	10.13	1.16	9.1		2.22	10.8
			333.7	10.50	1.16	9.1		2.22	10.8
			334.3	10.87	1.21	9.3		2.22	10.9
			334.9	11.98	1.37	9.4		2.22	10.7
			336.1	12.72	1.44	9.5	2176.0	2.32	10.5
			337.2	13.46	1.44	9.7		2.32	10.5
			337.8	14.21	1.44	9.9		2.32	10.4
			338.4	14.39	1.44	10.0	1582.0	2.32	10.7
			339.0	14.58	1.44	10.1		2.32	10.4
			340.2	15.88	1.55	10.2		2.32	10.3
			341.3	18.29	1.75	10.7		2.32	10.3
			342.5	19.03	1.80	10.9	1670.0	2.62	10.1
			343.1	19.58				2.67	10.1

R/V ACANIA 30 /NR / 1 MAY 79 CHEMICAL MISCOSCALE(PPJISE VI)									
TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PD4 PPM	NOTE: RATIO NO3/PD4	ATD NG/L	CHL A MG/93	TEMP DEG C
1300	36 6.5	121 59.1	343.7	20.14	1.84	10.9		1.31	10.0
			344.2	21.25	1.91	11.1		1.14	9.9
			345.4	22.37	1.99	11.2		0.96	9.8
			346.0	22.55	2.01	11.3		0.63	9.8
			346.6	22.74	2.01	11.2		0.47	9.8
			347.2	22.92	2.00	11.2		0.32	9.8
			347.7	22.37	2.00	11.2		0.50	9.8
			348.3	22.55	2.02	11.2		0.51	9.8
			348.9	22.55	2.02	11.2		0.70	9.8
			349.5	22.55	2.03	11.1		0.73	9.5
			350.1	22.55	2.04	11.2		0.72	9.5
			350.7	22.55	2.04	11.1		0.72	9.5
			351.2	22.92	2.05	11.2		0.75	9.5
			351.8	22.92	2.04	11.2		0.54	9.4
			352.4	22.92	2.04	11.2		0.70	9.4
			353.0	22.92	2.05	11.2		0.68	9.4
			353.6	22.92	2.06	11.2		0.66	9.4
			354.2	22.92	2.04	11.1		0.64	9.4
			354.7	22.92	2.03	11.1		0.63	9.4
			355.3	22.55	2.03	11.1		0.65	9.3
			355.9	22.00	2.00	11.0		0.51	9.7
			356.5	22.18	1.99	11.1		0.53	9.7
			357.1	21.81	1.93	11.0		0.55	9.7
			357.7	21.62	1.97	11.0		0.57	9.7
			358.2	20.33	1.91	10.6		0.46	9.9
			358.8	20.33	1.91	10.6		0.54	9.9
			359.4	20.33	1.91	10.6		0.31	10.0
			360.0	20.14	1.91	10.5		0.35	10.0
			360.6	22.33	1.91	11.7		0.39	10.0
			361.2	22.51	1.93	10.6			10.0
			361.7	20.70	1.94	10.8			10.0
			362.3	21.07	1.95	11.1			9.8
			362.9	21.25	1.91	11.1			9.8
			363.5	21.44	1.96	10.9			9.8
			364.1	21.44	1.96	10.9			9.7
			364.7	21.25	1.97	10.8			9.7
			365.2	20.88	1.97	10.6			9.7
			365.8	20.70	1.97	10.5			9.7
			366.4	20.70	1.97	10.5			9.7
1400	36 13.6	121 52.7					673.0	0.44	

R/V ACAN 1A

13 JUNE 79

CHEMICAL W. SECALF (CRUST VIT)

TIME GMT 2000	LATITUDE NORTH 36 36.3	LONGITUDE WEST 121 58.7	DISTANCE KM 0.4	NU ₃ UP 3.83	PO ₄ UM 0.73	NUIT. WATFIO M ³ /PO ₄ 5.2	ATO NG/L 497.0	CHL KG/93 1.53	T ^W W M ³ /C 13.8
			0.9	3.57	0.50	5.1		1.53	13.8
			1.3	3.54	0.50	7.7		1.53	13.8
			1.8	4.19	0.51	8.2		1.57	13.9
			2.2	6.18	0.69	9.1	1028.0	1.59	14.7
			2.6	8.16	0.79	10.3		1.57	14.3
			3.1	8.27	0.77	10.7		1.62	13.2
			3.5	8.45	0.77	11.0		1.63	13.4
			3.9	7.98	0.75	11.1		1.64	12.8
			4.4	7.62	0.70	10.9		1.67	12.8
			4.9	7.62	0.67	11.2		1.70	12.9
			5.3	6.47	0.58	11.4		1.72	12.9
			5.7	7.14	0.64	11.6	880.0	1.72	13.0
			6.1	7.98	0.66	12.1		1.73	12.8
			6.6	8.09	0.67	12.1		1.76	12.8
			7.0	7.62	0.65	11.7		1.77	12.7
			7.4	7.33	0.62	11.8		1.82	12.7
			7.9	9.08	0.60	11.8		1.82	12.8
			8.3					1.85	12.8
			8.8					1.85	12.9
			9.2					1.87	12.9
			9.6					1.89	12.9
			10.1					1.92	12.9
			10.5				1084.0	1.92	12.6
			11.0					1.93	12.4
			11.4					1.95	12.3
			11.8					1.96	12.3
			12.3					1.97	12.2
			12.7					1.99	12.2
			13.1					2.00	12.2
			13.7					2.02	11.6
			14.2					2.03	11.6
			14.8					2.05	11.6
			15.3					2.07	11.3
			15.9					2.07	11.3
			16.5					2.09	11.4
			17.0					2.10	11.4
			17.6					2.13	11.4
			18.1					2.15	11.4
			18.7					2.16	11.4
			19.2					2.18	11.4
			19.8					2.19	11.4
			20.3					2.20	11.4
			20.9					2.21	11.4
			21.5					2.22	11.4
			22.0					2.23	11.4
			22.6					2.24	11.4
			23.1					2.25	11.4
			23.7					2.26	11.4
			24.2					2.27	11.4

2100

36 29.2

121 57.1

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NCS °N	PO4 µM	NUTR. RATIO NH3/PO4	ATP NG/L	CHL. A NG/M3	TEMP °C
2200	36 20.4	121 57.1	24.8	11.05	1.14	9.7	1276.0	11.67	11.5
			29.3	11.23	1.18	9.5		11.63	11.5
			30.4	11.31	1.02	11.1		11.55	11.4
			31.0	11.45	1.03	11.1		11.53	11.4
			32.2	11.41	1.04	11.2	916.0	11.49	11.4
			32.8	11.63	1.10	10.8		11.45	11.4
			33.5	11.85	1.29	9.4		11.41	11.3
			34.1	12.10	1.29	9.3		11.39	11.3
			35.3	12.09	1.29	9.2	1276.0	11.37	11.3
			35.9	11.92	1.29	8.5		11.36	11.4
			36.5	11.74	1.22	8.5		11.32	11.4
			37.1	11.40	1.27	8.6		11.30	11.5
			38.4	11.41	1.24	8.6		11.29	11.5
			39.0	10.87	1.24	8.5		11.27	11.5
			39.6	10.51	1.22	8.4	1175.0	11.25	11.6
			40.2	10.19				11.23	11.6
			40.8					11.22	11.7
			41.4				825.0	11.20	11.7
			42.1					11.18	11.8
			42.7					11.15	11.9
			43.3					11.11	11.9
			43.9					11.07	12.0
			44.5	8.53	1.24	6.9		11.05	12.0
			45.1	8.60	1.18	7.3	1897.0	10.99	12.0
			45.7	8.78	1.15	7.6		10.93	12.0
			46.4	8.35	1.10	7.7		10.80	12.0
			47.0	8.13	1.06	8.0	1761.0	10.76	12.0
			48.2	7.98	1.00	8.1		10.71	12.0
			48.4	7.62	0.95	8.1		10.66	12.0
			48.6	7.30	0.90	8.1		10.61	12.0
			48.8	7.48	0.91	8.2		10.59	12.0
			49.0	7.26	0.88	7.8		10.52	12.0
			49.3	6.93	0.79	7.1	2057.0	10.48	12.0
			49.5	5.30	0.73	6.6		10.43	12.0
			49.7	4.30	0.65	6.6		10.37	12.0
			50.1	3.80	0.61	6.2		10.30	12.0
			50.3	2.97	0.52	5.7		10.27	12.0
2300	36 11.0	121 52.6							

R/V ACANIA

13 JUNE 79

CHEMICAL MESOSCALE(CRUISE VII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	UNC3	UNC4	NOTE-RATIO UNC3/UNC4	ATP NG/L	CUL MG/ML	TEMP C
			50.5	2.06	0.46	4.5	1857.0	11.31	13.2
			50.7	1.77	0.43	4.1		8.75	13.2
			50.9	1.77	0.43	4.1		4.30	13.4
			51.1	1.34	0.39	3.4		4.26	13.5
			51.4	0.58	0.36	2.7	2374.0	4.69	13.5
			51.6	0.87	0.34	2.6		2.13	13.5
			51.8	0.80	0.35	2.3		5.32	13.5
			52.0	0.76	0.35	2.2		5.57	13.5
			52.2	0.69	0.34	2.0		10.24	13.6
			52.4	0.55	0.33	1.7	1977.0	14.72	13.6
			52.6	0.66	0.33	2.0		14.93	13.6
			52.8	0.76	0.37	2.1		15.36	13.6
			53.0	0.87	0.41	2.2			13.6
			53.2	1.52	0.45	3.7			13.6
			53.5	2.21	0.45	5.0			13.6
			53.7	3.15	0.51	6.3	1298.0		13.6
			54.1	3.83	0.56	6.6			13.6
			54.3	4.05	0.56	6.8			13.6
			54.5	4.20	0.62	6.5	1851.0		12.7
			54.8	4.66	0.63	7.4			12.7
			55.1	4.55	0.62	7.3			12.7
			55.3	4.48	0.63	7.1			12.8
			55.6	5.67	0.70	8.1			12.8
			55.9	5.73	0.65	8.7	1421.0		12.9
			56.2	5.57	0.70	8.0			12.9
			56.5	5.02	0.67	7.5			12.9
			57.0	4.66	0.64	7.4			12.9
			57.3	4.59					12.9
			57.6						12.9
			57.8						12.9
			58.1						12.9
			58.4						12.9
			58.7						12.7
			59.0				1851.0		12.7
			59.2						12.6
			59.5						12.6
			59.8						12.6
			60.1						12.5
			60.4						12.5
			60.6						12.4
			60.9						12.4
			61.2						12.4
			61.5	5.02	0.49	10.2			12.3
			61.7	7.80	0.70	11.1			12.3
			62.0	8.09	0.74	10.9			12.3
			62.3	8.35	0.78	10.7			12.3
			62.6	8.71	0.82	10.6			12.1
			62.9						12.0

[illegible]

R/V ACANIA		13 JUNE 79		CHEMICAL MESOSCALE-CRUISE VII)					
TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	MUTP-RATIO NO3/PO4	ATP MG/L	CHL. a MG/M3	T-TEMP DEG C
0248	36	21.8	122 1.9	80.2	1.14	10.3			11.7
			80.4	11.23	1.00	11.2			11.7
			80.6	11.23	0.99	11.5			11.7
			80.8	11.52	1.17	9.8			11.7
			80.9	11.13	1.10	10.1			11.7
			81.1						11.7
			81.3						11.7
			81.5						11.7
			81.7						11.8
			81.9						11.8
			82.4						11.8
			83.4						11.9
			83.9						11.6
			84.5						11.7
			85.0						11.7

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR. PARTIO NO3/PO4	ATP NG/L	CHL A MG/M3	TWMP DEG C
0200	36 34.0	122 2.1	0.6					5.32	1.5
			31.7	1.80	0.99	1.8		5.13	1.3
			31.1	0.73	0.0	0.0		6.16	1.3
			29.8	0.37	0.0	0.0		6.06	1.3
			29.2	0.37	0.0	0.0		6.10	1.3
			27.9	0.55	0.94	0.6		6.40	1.3
			26.6					6.90	1.3
			25.0					7.19	1.3
			24.7					7.45	1.3
			24.1	1.71	0.92	1.9		7.52	1.3
			23.5	1.62		1.7		6.74	1.3
			22.2	3.89	0.93	3.4		6.45	1.3
			21.6	3.05	0.90	3.4		6.44	1.3
			20.9	3.89	0.93	3.4		5.27	1.3
			19.7	3.14	0.89	3.5		5.19	1.3
			18.4	5.46	0.91	5.3		5.27	1.3
			17.8	4.84	0.94	5.8		4.93	1.3
			17.1	0.0	0.03	0.0		4.44	1.3
			16.5	7.25	0.94	7.7		4.44	1.3
			15.9	3.30	0.95	4.8		4.44	1.3
			15.2	3.56	0.85	4.7		4.40	1.3
			14.6	4.30	0.84	5.1		3.94	1.3
			14.3	4.14	0.84	4.9		3.94	1.3
			13.9	4.12	0.84	4.9		4.11	1.3
			13.3	4.93	0.82	4.8		3.73	1.3
			12.7	3.91	0.82	4.8		3.73	1.3
			12.0	3.86	0.82	4.8		3.60	1.3
			11.4	4.12	0.82	5.0		3.63	1.3
			10.8	3.5	0.82	4.9		3.63	1.3
			10.1	3.87	0.82	4.7		3.63	1.3
			8.9	3.89	0.82	4.7		3.43	1.3
			8.22	8.22	0.82	4.7		3.43	1.3
			7.7	7.6	0.82	4.7		4.18	1.3
			6.3	6.3	0.82	4.7		4.18	1.3
			5.7	5.7	0.82	4.7		4.18	1.3
			5.1	5.1	0.82	4.7		4.18	1.3
			4.4	4.4	0.82	4.7		4.18	1.3
			3.8	3.8	0.82	4.7		4.18	1.3
			3.2	3.2	0.82	4.7		4.18	1.3
			2.5	2.5	0.82	4.7		4.18	1.3
			1.3	1.3	0.82	4.7		4.18	1.3
			0.6	0.6	0.82	4.7		4.18	1.3

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	PM3 UM	PM4 UM	NUTR. NO3/PM4	ATP NG/L	CHL A MG/ML	TEMP C
0300	36 26.7	122 9.9	32.3	1.41	0.0	0.0		5.05	13.7
			33.0	1.27	0.0	0.0		5.29	13.7
			33.6	1.73	1.04	1.7		4.45	13.7
			34.2					4.56	13.7
			35.5					3.32	13.7
			36.1	2.52	0.93	2.7		3.36	13.7
			37.4	1.77	0.93	1.9		3.78	13.7
			38.0	0.27	1.05	1.2		4.14	13.7
			39.7	0.20	1.21	0.5		4.13	13.7
			39.9	1.27	1.19	0.2		3.96	13.7
			40.6	1.62	1.25	0.8		4.02	13.7
			41.2	1.80	1.12	1.6		4.16	13.7
			41.8					4.17	13.7
			42.5					4.16	13.7
			43.1	8.52	0.92	9.3		4.06	13.7
			43.7	8.05	0.92	8.7		4.35	13.7
			44.4	8.59	0.92	9.3		4.13	13.7
			45.0	10.20	0.95	10.7		5.03	13.7
			45.6	10.66	1.01	10.6		5.34	13.7
			46.3	18.59	1.15	16.2		5.95	13.7
			47.6	20.91	1.23	17.0		5.94	13.7
			48.2	0.0	1.25	0.0		4.02	13.7
			49.8					3.94	13.7
			50.1					4.47	13.7
			50.7					4.41	13.7
			51.4	18.20	0.0	0.0		3.53	13.7
			52.0	18.23	0.0	0.0		3.67	13.7
			53.6		0.0	0.0		3.56	13.7
			53.9					3.69	13.7
			54.5					2.88	13.7
			55.2					2.33	13.7
			55.8					2.01	13.7
			56.4					1.84	13.7
			57.1					1.70	13.7
			57.7					1.71	13.7
			58.7					1.69	13.7
			59.0					1.77	13.7
			60.6					1.41	13.7
			61.5					1.27	13.7
			62.1					1.17	13.7
			62.8					1.13	13.7
			63.4					1.01	13.7
0400	36 19.2	122 17.6						0.95	13.3

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PD4 UM	NUTR. RATIO NO3/PD4	ATP NG/L	CHL A MG/L	TEMP DEG C
0500	36	10.0	122 25.7						
			64.0					0.54	13.3
			64.7					0.84	13.3
			65.3					0.85	13.5
			65.9					0.80	13.5
			66.6					0.79	13.6
			67.2					0.75	13.7
			67.8					0.76	13.9
			68.3					0.80	14.0
			68.9					0.79	14.0
			69.5					0.74	14.1
			70.0					0.74	14.1
			70.6					0.82	14.1
			71.2					0.89	14.1
			71.7					0.98	14.0
			72.3					0.97	14.0
			72.9					0.95	14.0
			73.4					0.95	14.0
			74.0					0.98	14.0
			74.6					1.02	14.1
			75.1					1.03	14.1
			75.7					1.05	14.1
			76.2					1.05	14.1
			76.8					1.05	14.1
			77.4					1.04	14.1
			77.9					0.99	14.1
			78.5					0.97	14.1
			79.1					0.99	14.0
			79.6					0.98	14.0
			80.2					0.99	14.0
			80.8					0.95	14.1
			81.3					0.95	14.1
			81.9					0.92	14.1
			82.5					0.93	14.0
			83.0					0.97	14.0
			83.6					0.97	14.0
			84.2					0.97	14.0
			84.7					0.97	14.0
			85.3					0.97	14.0
			85.9					0.97	14.0
			86.4					0.97	14.0
			87.0					0.97	14.0
			87.6					0.96	14.0
			88.1					0.97	14.0
			88.7					0.93	14.0
			89.3					0.90	14.0
			89.8					0.90	14.0
			90.4					0.89	14.0
			91.0					0.89	14.0
			91.5					0.85	14.0
			92.1					0.85	14.0
				9.12	1.17	7.8			
				9.40	1.07	8.8			
				9.46	0.99	7.6			
				9.86	1.27	7.8			
				9.67	1.21	8.0			

TIME GMT 0600	LATITUDE NORTH 36 2.1	LONGITUDE WEST 122 32.0	DISTANCE KM	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL. A MG/M ³	TEMP DEG C
			92.6					0.96	13.9
			92.2	8.15	0.97	8.4		0.96	13.9
			92.8	7.78	0.77	10.1		0.93	13.5
			94.3					0.93	13.5
			94.9					0.93	13.5
			95.5					0.92	13.5
			96.0	7.74	0.92	8.4		0.92	13.5
			96.6	7.67	0.93	8.2		0.95	13.9
			97.2	8.17	0.97	8.4		0.95	13.9
			97.7	8.16	1.21	6.7		0.91	13.5
			98.3	8.28	2.07	4.0		0.83	13.9
			93.9					0.81	13.9
			99.4					0.79	13.9
			100.0					0.87	13.9
			100.6					0.94	13.9
			101.1					0.98	13.5
			101.7					1.03	13.5
			102.3					1.06	14.0
			102.8					1.01	13.9
			103.4					1.01	13.9
			103.9					1.02	13.9
			104.5					1.00	13.9
			105.1					0.99	13.9
			105.6					0.99	13.9
			106.2					0.96	13.9
			106.7					0.95	13.9
			107.3					0.95	13.5
			107.9					0.91	13.5
			109.5					0.91	14.0
			109.0					0.92	14.4
			109.6					0.89	14.0
			110.2					0.89	14.0
			110.7					0.85	14.0
			111.3					0.85	15.0
			111.9					0.85	15.0
			112.4					0.85	15.3
			113.0					0.85	15.3
			113.6					0.86	15.4
			114.1					0.86	15.4
			114.7					0.86	15.4
			115.3					0.86	15.4
			115.8					0.85	15.4
			116.4					0.85	15.4
			117.0					0.85	15.4
			117.5					0.85	15.4
			118.1					0.87	15.3
			118.6					0.87	15.3
			119.2					0.94	15.3
			119.8					0.95	15.4
			120.3					0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87	15.4
								0.87	15.4
								0.94	15.3
								0.95	15.4
								0.95	15.4
								0.85	15.4
								0.83	15.4
								0.87</	

R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE(CRUISE VITI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NT3 UM	PO4 UM	NUTR.RATIO NU3/PO4	ATD NG/L	CHI A MG/M3	TEMP C
			149.6	7.52	0.0	0.0		C.52	14.6
			150.1	8.18	0.0	0.0		C.55	14.5
			150.6	8.70	0.0	0.0		C.53	14.5
			151.1	8.25	0.0	0.0		C.74	14.5
			151.6	8.77	0.0	0.0		C.71	14.5
			152.0	8.96	0.0	0.0		C.69	14.5
			152.5	8.70	0.0	0.0		C.72	14.5
			153.0	7.66	0.0	0.0		C.73	14.5
			153.5	7.35	0.0	0.0		C.74	14.5
			154.0	7.14	0.0	0.0		C.71	15.1
			154.5	7.14	0.0	0.0		C.69	15.1
			155.0	7.09	0.0	0.0		C.69	15.1
			155.5	7.27	0.0	0.0		C.67	15.1
			156.0	7.32	0.0	0.0		C.66	15.1
			156.4					C.63	15.0
			157.0					C.62	14.9
			157.4					C.61	14.8
			157.9					C.59	14.7
			158.4					C.55	14.7
			158.9					C.52	14.5
			159.4					C.50	14.5
			160.0					C.47	14.5
			160.3					C.44	14.5
			160.8					C.44	14.5
			161.3					C.44	14.5
			161.8					C.44	14.5
			162.3					C.44	14.5
			162.8					C.44	14.5
			163.3					C.44	14.5
			163.8					C.44	14.5
			164.3					C.44	14.5
			164.8					C.44	14.5
			165.3					C.44	14.5
			165.8					C.44	14.5
			166.3					C.44	14.5
			166.8					C.44	14.5
			167.3					C.44	14.5
			167.8					C.44	14.5
			168.3					C.44	14.5
			168.8					C.44	14.5
			169.3					C.44	14.5
			169.8					C.44	14.5
			170.3					C.44	14.5
			170.8					C.44	14.5
			171.3					C.44	14.5
			171.8					C.44	14.5
			172.3					C.44	14.5
			172.8					C.44	14.5
			173.3					C.44	14.5
			173.8					C.44	14.5
			174.3					C.44	14.5

1000 35 29.1 122 57.1

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PD4 UM	NUP:RATIO NO3/PD4	ATP NG/L	CHL A MG/L	TEMP C
0800	35 42.0	122 44.8	125.6	3.03	0.95	3.2	0.84	1.09	15.3
			127.2	3.17	0.89	3.6	0.84	1.04	15.3
			129.0	3.38	0.89	3.8	0.84	1.04	15.3
			129.6	3.07	0.87	3.5	0.84	0.96	15.3
			130.3	3.36	0.89	3.8	0.84	0.94	15.3
			131.5	3.41	0.89	3.7	0.84	0.94	15.3
			132.1	3.54	0.89	3.8	0.84	0.94	15.3
			133.3						15.3
			133.9						15.3
			134.5	4.33	0.75	5.8	0.68	0.70	15.1
			135.1	4.14	0.71	5.1	0.68	0.65	15.1
			136.3	4.26	0.78	5.5	0.68	0.62	15.2
			137.5	4.33	0.85	5.3	0.68	0.64	15.2
			138.7	4.54	0.82	5.6	0.68	0.65	15.1
			139.3	4.77	0.79	6.1	0.66	0.66	15.1
			140.5	4.86	0.79	6.4	0.66	0.66	15.1
			141.1						15.0
			141.7						15.0
			142.3						15.0
			142.9						15.0
			143.5	3.66	0.94	3.9	0.85	0.85	15.0
			144.1	6.13	0.90	6.9	0.85	0.85	15.0
			145.2	6.36	0.92	6.9	0.85	0.85	15.0
			145.7	6.41	0.98	7.7	0.85	0.85	14.9
			146.2	6.70	0.91	7.5	0.85	0.85	14.9
			146.7	7.14	0.96	7.4	0.83	0.84	14.9
			147.2	7.40	0.0	0.0	0.83	0.83	14.8
			147.6	7.89	0.0	0.0	0.82	0.82	14.7
			148.1	8.12	0.0	0.0	0.82	0.82	14.7
			148.6	7.92	0.0	0.0	0.82	0.82	14.7
			149.1						14.7
0900	35 36.4	122 49.9	144.1	3.66	0.94	3.9	0.85	0.85	15.0
			144.7	6.13	0.90	6.9	0.85	0.85	15.0
			145.2	6.36	0.92	6.9	0.85	0.85	15.0
			145.7	6.41	0.98	7.7	0.85	0.85	14.9
			146.2	6.70	0.91	7.5	0.85	0.85	14.9
			146.7	7.14	0.96	7.4	0.83	0.84	14.9
			147.2	7.40	0.0	0.0	0.83	0.83	14.8
			147.6	7.89	0.0	0.0	0.82	0.82	14.7
			148.1	8.12	0.0	0.0	0.82	0.82	14.7
			148.6	7.92	0.0	0.0	0.82	0.82	14.7
			149.1						14.7

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE NM	NC3 UN	PD4 HPI	NUTP-RATIO NO3/PD4	ATP NG/L	CHL MG/43	TEMP C
1452'	35 47.3	123 24.7	234.5	P.02	7.65	9.44		1.15	15.5
			235.2	8.02	7.65	9.44		1.24	15.5
			235.8	7.87	0.89	8.9		0.52	15.5
			236.4	7.73	0.80	9.7		1.24	15.5
			237.1	7.59	0.85	8.9		1.15	15.5
			237.7	7.44	0.85	8.9		1.09	15.5
			238.3	7.73	0.88	8.8		1.08	15.5
			239.0	7.99	0.87	9.2		1.06	15.5
			239.6	7.10	0.81	8.8		1.04	15.5
			240.2	7.87	0.87	9.0		0.57	15.5
			240.9	7.96	0.88	9.0		0.57	15.5
			241.5	8.02	0.88	9.1		0.52	15.5
			242.2	8.13	0.83	9.2		1.00	15.5
			242.9	7.73	0.77	9.0		0.95	15.5
			243.4	8.39	0.87	9.6		0.51	15.5
			244.1	8.53	0.90	9.5		0.89	15.5
			244.7	8.68	0.83	9.8		0.89	15.5
			245.3	8.73	0.83	9.9		0.89	15.5
			246.0	8.30	0.0	9.0		0.89	15.5
			246.6	8.82	0.0	9.0		0.89	15.5
			247.2	9.25	1.01	9.2		0.89	15.5
			247.9						15.5
			249.1						15.5
			250.4						15.5
			251.0						15.5
			251.7	10.01	1.01	9.9		0.32	15.5
			252.3	10.25	1.00	10.3		0.35	15.5
			252.9	7.99	0.85	9.4		0.36	15.5
			254.2	6.23	0.78	7.3		0.38	15.5
			254.8	5.67	0.74	7.6		0.40	15.5
			255.5	5.25	0.74	6.9		0.41	15.5
			256.1	5.10	0.76	6.8		0.48	15.5
			257.4	5.44	0.78	7.0		0.49	15.5
			258.0	5.85	0.81	7.5		0.49	15.5
			259.3	6.21	0.83	7.7		0.49	15.5
			260.5	6.63	0.83	8.0		0.49	15.5
			261.2	7.01	0.83	8.4		0.48	15.5
			261.8	7.43	0.83	8.8		0.48	15.5
			262.4	7.86	0.83	9.1		0.48	15.5
			263.1	8.28	0.83	9.4		0.48	15.5
			263.7	8.70	0.83	9.7		0.48	15.5
			264.3	9.12	0.83	10.0		0.48	15.5
			265.0	9.54	0.83	10.3		0.48	15.5
			265.6	9.96	0.83	10.6		0.48	15.5

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE FM	NR3 UM	PT4 UM	NR3/PT4	ATP NG/L	CHL A MG/M2	TEMP DEG C
1600	35	36.5	123	266.2	0.04	0.02	8.1	0.44	16.1
			266.9	4.80	0.66	7.3	0.43	0.43	16.1
			267.5	4.58	0.0	0.0	0.0	0.40	16.2
			268.1	4.56	0.55	8.1	0.38	0.38	16.4
			269.4	4.46	0.55	8.3	0.36	0.36	16.5
			270.0	4.39	0.53	8.3	0.33	0.33	16.5
			270.7	4.46	0.52	8.6	0.31	0.31	16.8
			271.3	4.36	0.52	8.2	0.29	0.29	16.5
			271.9	4.28	0.53	8.2	0.27	0.27	17.0
			272.6	4.21	0.53	7.9	0.25	0.25	17.0
			273.2	4.19	0.53	7.9	0.23	0.23	17.0
			273.8	4.24	0.51	8.3	0.23	0.23	16.8
			274.5				0.21	0.21	16.5
			275.1				0.19	0.19	16.9
			275.7				0.17	0.17	16.5
			276.4				0.17	0.17	16.9
			277.0				0.17	0.17	17.0
			277.6				0.17	0.17	17.0
			278.3				0.17	0.17	17.1
			279.9				0.17	0.17	17.1
			280.2				0.17	0.17	17.1
			280.8				0.17	0.17	17.1
			281.5				0.17	0.17	17.1
			282.1				0.17	0.17	17.1
			282.7				0.17	0.17	17.1
			283.4				0.17	0.17	17.1
			284.0				0.17	0.17	17.1
			284.6				0.17	0.17	17.1
			285.3				0.19	0.19	17.1
			285.9				0.18	0.18	17.1
			286.5				0.19	0.19	17.1
			287.2				0.19	0.19	17.1
			287.8				0.19	0.19	17.1
			289.4				0.18	0.18	17.1
			289.1				0.18	0.18	17.1
			290.3				0.17	0.17	17.1
			291.0				0.17	0.17	17.1
			291.6				0.17	0.17	17.1
			292.2				0.17	0.17	17.1
			292.9				0.17	0.17	17.1
			293.5				0.17	0.17	17.1
			294.1				0.17	0.17	17.1
			294.8				0.17	0.17	17.1
			295.4				0.17	0.17	17.1
			296.0				0.17	0.17	17.1
			296.7				0.18	0.18	17.1
			297.3				0.17	0.17	17.1

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UM	PD4 UM	MUTP-RATIO NO3/PO4	ATP MG/L	CHL A MG/43	TEMP DEG C
1805	34 32.8	123 32.3	297.0	0.0	0.25	0.0	188.0	0.17	17.1
			298.0	0.0	0.25	0.0		0.17	17.1
			299.0	0.0	0.25	0.0		0.17	17.1
			300.0	0.0	0.25	0.0		0.17	17.1
			301.0	1.09	0.25	4.00		0.17	17.1
			302.0	0.87	0.25	3.06		0.17	17.1
			303.0	0.0	0.25	4.00		0.17	17.1
			304.0	0.89	0.27	3.03		0.17	17.0
			305.0	0.89	0.27	3.03		0.17	17.0
			306.0	0.99	0.27	3.03		0.17	17.0
			307.0	0.92	0.27	3.03		0.17	17.0
			308.0	1.08	0.27	3.03		0.17	17.0
			309.0	0.98	0.24	3.03		0.19	17.0
			310.0	0.95	0.29	3.03		0.20	17.0
			311.0	0.96	0.25	3.03		0.21	17.0
			312.0	1.13	0.27	4.04	925.0	0.21	16.9
			313.0	1.00	0.26	3.08		0.22	16.9
			314.0	0.53	0.26	3.08		0.26	16.5
			315.0	0.99	0.26	3.08		0.28	16.5
			316.0	0.93	0.26	3.07	1165.0	0.29	16.6
			317.0	0.93	0.25	3.07		0.29	16.6
			318.0	0.93	0.25	3.07		0.29	16.6
			319.0	0.83	0.25	3.07		0.33	16.6
			320.0	0.89	0.25	3.06		0.34	16.6
			321.0	1.02	0.26	3.07		0.37	16.6
			322.0	0.95	0.26	4.01	1014.0	0.39	16.7
			323.0	1.22	0.32	3.09		0.40	16.6
			324.0	1.31	0.32	3.09		0.41	16.6
			325.0	8.10	0.35	25.09		0.44	16.6
			326.0	4.16	0.35	11.09		0.44	16.6
			327.0	0.45	0.39	13.05		0.47	16.6
			328.0	6.06	0.45		3418.0	0.47	16.5
			329.0					0.47	16.5
1855	34 39.4	123 36.0	321.0					0.46	16.5
			322.0					0.46	16.5
			323.0					0.46	16.4
			324.0					0.46	16.4
			325.0					0.46	16.4
			326.0					0.46	16.4
			327.0					0.46	16.4
			328.0					0.46	16.4
			329.0					0.46	16.4
1900	34 39.9	123 36.4	330.0					0.46	16.4
			331.0					0.46	16.4
			332.0					0.46	16.4
			333.0					0.46	16.4
			334.0					0.46	16.4
			335.0					0.46	16.4
			336.0					0.46	16.4
			337.0					0.46	16.4
			338.0					0.46	16.4
			339.0					0.46	16.4
			340.0					0.46	16.4
			341.0					0.46	16.4
			342.0					0.46	16.4
			343.0					0.46	16.4
			344.0					0.46	16.4
			345.0					0.46	16.4
			346.0					0.46	16.4
			347.0					0.46	16.4
			348.0					0.46	16.4
			349.0					0.46	16.4
			350.0					0.46	16.4
			351.0					0.46	16.4
			352.0					0.46	16.4
			353.0					0.46	16.4
			354.0					0.46	16.4
			355.0					0.46	16.4
			356.0					0.46	16.4
			357.0					0.46	16.4
			358.0					0.46	16.4
			359.0					0.46	16.4
			360.0					0.46	16.4
			361.0					0.46	16.4
			362.0					0.46	16.4
			363.0					0.46	16.4
			364.0					0.46	16.4
			365.0					0.46	16.4
			366.0					0.46	16.4
			367.0					0.46	16.4
			368.0					0.46	16.4
			369.0					0.46	16.4
			370.0					0.46	16.4
			371.0					0.46	16.4
			372.0					0.46	16.4
			373.0					0.46	16.4
			374.0					0.46	16.4
			375.0					0.46	16.4
			376.0					0.46	16.4
			377.0					0.46	16.4
			378.0					0.46	16.4
			379.0					0.46	16.4
			380.0					0.46	16.4
			381.0					0.46	16.4
			382.0					0.46	16.4
			383.0					0.46	16.4
			384.0					0.46	16.4
			385.0					0.46	16.4
			386.0					0.46	16.4
			387.0					0.46	16.4
			388.0					0.46	16.4
			389.0					0.46	16.4
			390.0					0.46	16.4
			391.0					0.46	16.4
			392.0					0.46	16.4
			393.0					0.46	16.4
			394.0					0.46	16.4
			395.0					0.46	16.4
			396.0					0.46	16.4
			397.0					0.46	16.4
			398.0					0.46	16.4
			399.0					0.46	16.4
			400.0					0.46	16.4
			401.0					0.46	16.4
			402.0					0.46	16.4
			403.0					0.46	16.4
			404.0					0.46	16.4
			405.0					0.46	16.4
			406.0					0.46	16.4
			407.0					0.46	16.4
			408.0					0.46	16.4
			409.0					0.46	16.4
			410.0					0.46	16.4
			411.0					0.46	16.4
			412.0					0.46	16.4
			413.0					0.46	16.4
			414.0					0.46	16.4
			415.0					0.46	16.4
			416.0					0.46	16.4
			417.0					0.46	16.4
			418.0					0.46	16.4
			419.0					0.46	16.4
			420.0					0.46	16.4
			421.0					0.46	16.4
			422.0					0.46	16.4
			423.0					0.46	16.4
			424.0					0.46	16.4
			425.0					0.46	16.4
			426.0					0.46	16.4
			427.0					0.46	16.4
			428.0					0.46	16.4
			429.0					0.46	16.4
			430.0					0.46	16.4
			431.0					0.46	16.4
			432.0					0.46	16.4
			433.0					0.46	16.4
			434.0					0.46	16.4
			435.0					0.46	16.4
			436.0					0.46	16.4
			437.0					0.46	16.4
			438.0					0.46	16.4
			439.0					0.46	16.4
			440.0					0.46	16.4
			441.0					0.46	16.4
			442.0					0.46	16.4
			443.0					0.46	16.4
			444.0					0.46	16.4
			445.0					0.46	16.4
			446.0					0.46	16.4
			447.0					0.46	16.4
			448.0					0.46	16.4
			449.0					0.46	16.4
			450.0					0.46	16.4
			451.0					0.46	16.4
			452.0					0.46	16.4
			453.0					0.46	16.4
			454.0					0.46	16.4
			455.0					0.46	16.4
			456.0					0.46	16.4
			457.0					0.46	16.4
			458.0					0.46	16.4
			459.0					0.46	16.4
			460.0					0.46	16.4
			461.0					0.46	16.4
			462.0					0.46	16.4
			463.0					0.46	16.4
			464.0					0.46	16.4
			465.0					0.46	16.4
			466.0					0.46	16.4
			467.0					0.46	16.4
			468.0					0.46	16.4
			469.0					0.46	16.4
			470.0					0.46	16.4
			471.0					0.46	16.4
			472.0					0.46	16.4
			473.0					0.46	16.4

TIME GMT	LATITUDE NCRTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR-RATIO NO3/PO4	ATP NG/L	CHL A MG/43	TEMP DEG C
1931	34 43.9	123 39.4	329.6				1082.0	0.47	15.6
			330.3					1.02	15.6
			331.5					1.71	15.6
			332.2					0.70	15.7
			332.8					0.69	15.7
			333.4					0.75	15.7
			334.1				1113.0	1.02	15.7
			334.7					1.01	15.6
			335.3					0.69	15.6
			336.0					0.69	15.6
			336.6				983.0	0.67	15.6
			337.2					0.69	15.6
			337.9					0.69	15.6
			338.5					0.64	15.6
			339.1					0.67	15.6
			339.9				1047.0	0.67	15.6
			340.4					0.67	15.6
			341.0					0.61	15.6
			341.7					0.61	15.6
			342.4					0.59	15.6
			343.0				1013.0	0.59	15.4
			343.6					0.61	15.4
			344.2					0.61	15.4
			344.8					0.63	15.4
			345.5					0.63	15.4
			346.1				981.0	0.63	15.4
			346.7					0.57	15.4
			347.4					0.61	15.5
			348.0					0.59	15.5
			348.6					0.59	15.5
			349.3				964.0	0.58	15.5
			349.9					0.58	15.5
			350.6					0.58	15.5
			351.2					0.55	15.5
			351.8					0.62	15.5
			352.4				1056.0	0.60	15.5
			353.1					0.56	15.5
			353.7					0.50	15.5
			354.3					0.47	15.5
			355.0					0.47	15.5
			355.6					0.43	15.5
			356.2					0.43	15.5
			356.9				977.0	0.40	15.4
			357.5					0.40	15.4
			358.1					0.39	15.4
			358.8					0.37	17.0
			359.4					0.37	17.0
			360.0				723.0	0.36	17.0
			360.7					0.35	16.8
2000	34 42.2	123 42.6							

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	MC3 UM	PO4 UM	NOTE NO3/PO4	ATP MG/L	CHL A MG/M3	TEMP DEC C
2100	34 36.3	123 51.4	361.3	0.0	0.79	0.0		0.34	16.5
			361.9	0.0	0.50	0.0		0.32	16.5
			362.6	0.0	0.50	0.0		0.37	17.0
			363.2	0.0	0.52	0.0		0.39	17.0
			363.8	0.0	0.53	0.0		0.29	17.1
			364.5	0.0	0.53	0.0		0.27	17.1
			365.1	0.0	0.53	0.0		0.26	17.1
			365.7	0.0	0.53	0.0		0.25	17.1
			366.4	0.0	0.47	0.0		0.24	17.1
			367.0	0.0	0.50	0.0		0.23	17.2
			367.6	0.0	0.49	0.0		0.22	17.2
			368.3	0.0	0.49	0.0		0.20	17.2
			369.0	0.0	0.48	0.0		0.19	17.2
			369.5	0.0	0.48	0.0		0.19	17.2
2128	34 33.9	123 55.0	370.2	0.0	0.50	0.0		0.18	17.2
			371.4	0.0	0.51	0.0		0.20	17.2
			372.1	0.0	0.52	0.0		0.21	17.2
			372.7	0.0	0.53	0.0		0.23	17.2
			373.3	0.0	0.51	0.0		0.25	17.2
			374.0	0.0	0.51	0.0		0.27	17.2
			374.6	0.0	0.51	0.0		0.29	17.2
			375.3	0.0	0.51	0.0		0.31	17.2
			375.9	0.0	0.51	0.0		0.32	17.2
2145	34 35.5	123 54.0	376.5	0.0	0.51	0.0		0.34	17.2
			377.2	0.0	0.51	0.0		0.35	17.1
			377.8	0.0	0.51	0.0		0.35	17.1
			378.4	0.0	0.51	0.0		0.35	17.1
			379.1	0.0	0.51	0.0		0.35	17.0
2200	34 37.4	123 54.7	379.7	0.0	0.51	0.0		0.38	17.0
			380.3	0.0	0.51	0.0		0.38	16.5
			381.0	0.0	0.51	0.0		0.42	16.5
			381.6	0.0	0.51	0.0		0.44	16.5
			382.2	0.0	0.51	0.0		0.44	16.5
			382.9	0.0	0.51	0.0		0.44	15.5
			383.5	0.0	0.51	0.0		0.44	15.5
			384.1	0.0	0.51	0.0		0.44	15.5
			384.9	0.0	0.51	0.0		0.44	15.5
			385.4	0.0	0.51	0.0		0.44	15.5
			386.0	0.0	0.51	0.0		0.44	15.5
			386.7	0.0	0.51	0.0		0.44	15.5
			387.3	0.0	0.51	0.0		0.44	15.5
			387.9	0.0	0.51	0.0		0.44	15.5
			388.5	0.0	0.51	0.0		0.44	15.5
			389.2	0.0	0.51	0.0		0.44	15.5
			389.8	0.0	0.51	0.0		0.44	15.5
			390.5	0.0	0.51	0.0		0.44	15.5
			391.1	0.0	0.51	0.0		0.44	15.5
			391.7	0.0	0.51	0.0		0.44	15.5
			392.4	0.0	0.51	0.0		0.44	15.5
2232	34 42.0	123 54.1		0.0	0.51	0.0		0.44	15.5

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR. RATIO NO3:PO4	ATP NG/L	CHL. A MG/M3	TEMP. C
2300	34 44.4	123 51.6	393.0	3.34	0.74	4.5		0.48	15.3
			393.6	3.11	0.69	2.1		0.53	14.5
			394.3	1.35	0.63	1.2		0.51	14.9
			394.9	0.71	0.61	1.2		0.50	14.8
			395.5	0.79	0.64	1.2		0.49	14.7
			396.2	0.46	0.0	0.0	1188.0	0.49	14.7
			397.4					0.47	14.7
			398.1					0.43	14.5
			399.3				1098.0	0.43	14.5
			400.0					0.42	15.0
			400.6					0.42	15.0
			401.2					0.42	15.0
			401.7					0.44	15.1
			402.5					0.44	15.2
			403.1					0.44	15.2
2400	34 49.3	123 45.0	403.8					0.45	15.2
			404.4					0.44	15.2
			405.0					0.44	15.3
			405.7					0.44	15.3
			406.3					0.44	15.3
			406.9					0.44	15.3
			407.6				1276.0	0.45	15.3
			408.2					0.45	15.4
			408.8					0.45	15.4
			409.5					0.45	15.3
			410.1				1102.0	0.46	15.3
			410.7					0.46	15.3
			411.4					0.46	15.2
			412.0					0.46	15.1
			412.6					0.48	15.1
			413.3					0.49	15.0
			413.9					0.50	15.0
			414.5					0.50	15.1
			415.2					0.54	15.1
			415.9					0.54	14.9
			416.4					0.57	14.9
			417.1					0.54	14.8
			417.7					0.54	14.8
			419.3					0.52	14.6
			419.9					0.52	14.4
			419.6					0.52	14.4
			420.3					0.52	14.1
			420.9					0.52	14.0
			421.5					0.52	13.5
			422.2					0.54	13.8
			422.8					0.52	13.9
			423.4					0.52	13.9
			424.1					0.52	13.9

TIME GMT	LATITUDE NCRTH	LONGITUDE WEST	DISTANCE KM	NU3 UM	PD4 UM	NU3/PD4 RATIO	ATP NG/L	CHL A MG/L	TEMP DEG C
0100	34 55.4	123 38.1	424.7	2.03	0.70	2.9	0.52	0.52	14.0
			425.3	2.16	0.71	3.0	0.52	0.52	14.0
			426.0	2.07	0.71	2.9	0.52	0.52	14.0
			426.6	1.89	0.71	2.7	0.52	0.52	14.0
			427.2	1.21	0.69	1.8	0.52	0.52	14.0
			427.9				0.52	0.52	14.0
			428.5				0.52	0.52	14.0
			429.1				0.52	0.52	14.0
			429.8				0.52	0.52	14.0
			430.4				0.52	0.52	14.0
			431.0				0.52	0.52	14.0
			431.7				0.52	0.52	14.0
			432.3				0.52	0.52	14.0
			432.9				0.52	0.52	14.0
			433.6				0.52	0.52	14.0
			434.2				0.52	0.52	14.0
			434.8				0.52	0.52	14.0
			435.5				0.52	0.52	14.0
			436.1				0.52	0.52	14.0
			436.7				0.52	0.52	14.0
			437.4				0.52	0.52	14.0
			438.0				0.52	0.52	14.0
			438.6	1.72	0.79	2.2	0.52	0.52	14.0
			439.3	3.03	0.80	3.8	0.52	0.52	14.0
			440.0	2.64	0.78	3.4	0.52	0.52	14.0
			441.2	1.54	0.78	2.5	0.52	0.52	14.0
			441.8	3.69	0.80	4.6	0.52	0.52	14.0
			442.4	4.34	0.83	5.2	0.52	0.52	14.0
			443.1	6.09	0.87	7.0	0.52	0.52	14.0
			444.3	8.14	0.90	9.0	0.52	0.52	14.0
			444.8	5.87	0.92	6.5	0.52	0.52	14.0
			445.0	5.09	0.94	6.7	0.52	0.52	14.0
			445.6	5.32	0.87	6.3	0.52	0.52	14.0
			446.2	4.04	0.86	4.9	0.52	0.52	14.0
			447.5	4.78	0.87	5.5	0.52	0.52	14.0
			448.1	5.69	0.91	6.3	0.52	0.52	14.0
			448.9	6.74	0.92	7.3	0.52	0.52	14.0
			449.4	6.61	0.89	7.4	0.52	0.52	14.0
			450.0	5.61	0.87	6.9	0.52	0.52	14.0
			451.3	6.26	0.86	7.1	0.52	0.52	14.0
			452.6	6.04	0.85	7.0	0.52	0.52	14.0
			453.2	6.09	0.85	7.0	0.52	0.52	14.0
			453.8	5.69	0.85	6.7	0.52	0.52	14.0
			454.5	5.56	0.85	6.5	0.52	0.52	14.0
			455.1				0.52	0.52	14.0
			455.7				0.52	0.52	14.0

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE NM	UM _{NC3}	UM _{PD4}	NUTR. RATIO NM3/PO4	ATP NG/L	CHL A MG/ML	TEMP DEG C	
0200	35	1.9	123 31.7	5.65 5.61 5.74	0.85 0.85 0.83	6.6 6.6 6.9		0.75 0.77 0.86	14.5 15.0 14.9	
			456.4 457.0 457.6 458.3 459.9 459.5 460.2 460.9 461.4 462.1 462.7 463.3 464.0 464.6 465.2 465.9 466.5 467.2 467.8 468.4 469.1 470.3 471.0 471.6 472.2 472.9 473.5 474.1 474.8 475.4 476.0 476.7 477.3 477.9 478.6 479.2 479.8 480.5 481.1 481.7 482.4 483.0 483.6 484.3 484.9 485.5 486.2 486.8 487.4	5.34 5.73 5.21 5.39 5.69 5.82 6.74 6.43 6.22 5.17 4.25 2.42	0.87 0.88 0.85 0.83 0.85 0.85 0.85 0.85 0.85 0.84 0.80	6.1 5.4 6.1 6.5 6.9 6.8 7.9 7.6 7.3 6.2 5.3		0.73 0.77 0.77 0.77 0.77 0.77 0.78	1.08 1.17 1.20	14.5 15.0 14.9 14.6 14.5 14.4 14.3 14.2 14.1 14.0 13.9 13.8 13.7 13.6 13.5 13.4 13.3 13.2 13.1 13.0 12.9 12.8 12.7 12.6 12.5 12.4 12.3 12.2 12.1 12.0 11.9 11.8 11.7 11.6 11.5 11.4 11.3 11.2 11.1 11.0 10.9 10.8 10.7 10.6 10.5 10.4 10.3 10.2 10.1 10.0 9.9 9.8 9.7 9.6 9.5 9.4 9.3 9.2 9.1 9.0 8.9 8.8 8.7 8.6 8.5 8.4 8.3 8.2 8.1 8.0 7.9 7.8 7.7 7.6 7.5 7.4 7.3 7.2 7.1 7.0 6.9 6.8 6.7 6.6 6.5 6.4 6.3 6.2 6.1 6.0 5.9 5.8 5.7 5.6 5.5 5.4 5.3 5.2 5.1 5.0 4.9 4.8 4.7 4.6 4.5 4.4 4.3 4.2 4.1 4.0 3.9 3.8 3.7 3.6 3.5 3.4 3.3 3.2 3.1 3.0 2.9 2.8 2.7 2.6 2.5 2.4 2.3 2.2 2.1 2.0 1.9 1.8 1.7 1.6 1.5 1.4 1.3 1.2 1.1 1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2 0.1 0.0 -0.1 -0.2 -0.3 -0.4 -0.5 -0.6 -0.7 -0.8 -0.9 -1.0 -1.1 -1.2 -1.3 -1.4 -1.5 -1.6 -1.7 -1.8 -1.9 -2.0 -2.1 -2.2 -2.3 -2.4 -2.5 -2.6 -2.7 -2.8 -2.9 -3.0 -3.1 -3.2 -3.3 -3.4 -3.5 -3.6 -3.7 -3.8 -3.9 -4.0 -4.1 -4.2 -4.3 -4.4 -4.5 -4.6 -4.7 -4.8 -4.9 -5.0 -5.1 -5.2 -5.3 -5.4 -5.5 -5.6 -5.7 -5.8 -5.9 -6.0 -6.1 -6.2 -6.3 -6.4 -6.5 -6.6 -6.7 -6.8 -6.9 -7.0 -7.1 -7.2 -7.3 -7.4 -7.5 -7.6 -7.7 -7.8 -7.9 -8.0 -8.1 -8.2 -8.3 -8.4 -8.5 -8.6 -8.7 -8.8 -8.9 -9.0 -9.1 -9.2 -9.3 -9.4 -9.5 -9.6 -9.7 -9.8 -9.9 -10.0 -10.1 -10.2 -10.3 -10.4 -10.5 -10.6 -10.7 -10.8 -10.9 -11.0 -11.1 -11.2 -11.3 -11.4 -11.5 -11.6 -11.7 -11.8 -11.9 -12.0 -12.1 -12.2 -12.3 -12.4 -12.5 -12.6 -12.7 -12.8 -12.9 -13.0 -13.1 -13.2 -13.3 -13.4 -13.5 -13.6 -13.7 -13.8 -13.9 -14.0 -14.1 -14.2 -14.3 -14.4 -14.5 -14.6 -14.7 -14.8 -14.9 -15.0 -15.1 -15.2 -15.3 -15.4 -15.5 -15.6 -15.7 -15.8 -15.9 -16.0 -16.1 -16.2 -16.3 -16.4 -16.5 -16.6 -16.7 -16.8 -16.9 -17.0 -17.1 -17.2 -17.3 -17.4 -17.5 -17.6 -17.7 -17.8 -17.9 -18.0 -18.1 -18.2 -18.3 -18.4 -18.5 -18.6 -18.7 -18.8 -18.9 -19.0 -19.1 -19.2 -19.3 -19.4 -19.5 -19.6 -19.7 -19.8 -19.9 -20.0 -20.1 -20.2 -20.3 -20.4 -20.5 -20.6 -20.7 -20.8 -20.9 -21.0 -21.1 -21.2 -21.3 -21.4 -21.5 -21.6 -21.7 -21.8 -21.9 -22.0 -22.1 -22.2 -22.3 -22.4 -22.5 -22.6 -22.7 -22.8 -22.9 -23.0 -23.1 -23.2 -23.3 -23.4 -23.5 -23.6 -23.7 -23.8 -23.9 -24.0 -24.1 -24.2 -24.3 -24.4 -24.5 -24.6 -24.7 -24.8 -24.9 -25.0 -25.1 -25.2 -25.3 -25.4 -25.5 -25.6 -25.7 -25.8 -25.9 -26.0 -26.1 -26.2 -26.3 -26.4 -26.5 -26.6 -26.7 -26.8 -26.9 -27.0 -27.1 -27.2 -27.3 -27.4 -27.5 -27.6 -27.7 -27.8 -27.9 -28.0 -28.1 -28.2 -28.3 -28.4 -28.5 -28.6 -28.7 -28.8 -28.9 -29.0 -29.1 -29.2 -29.3 -29.4 -29.5 -29.6 -29.7 -29.8 -29.9 -30.0 -30.1 -30.2 -30.3 -30.4 -30.5 -30.6 -30.7 -30.8 -30.9 -31.0 -31.1 -31.2 -31.3 -31.4 -31.5 -31.6 -31.7 -31.8 -31.9 -32.0 -32.1 -32.2 -32.3 -32.4 -32.5 -32.6 -32.7 -32.8 -32.9 -33.0 -33.1 -33.2 -33.3 -33.4 -33.5 -33.6 -33.7 -33.8 -33.9 -34.0 -34.1 -34.2 -34.3 -34.4 -34.5 -34.6 -34.7 -34.8 -34.9 -35.0 -35.1 -35.2 -35.3 -35.4 -35.5 -35.6 -35.7 -35.8 -35.9 -36.0 -36.1 -36.2 -36.3 -36.4 -36.5 -36.6 -36.7 -36.8 -36.9 -37.0 -37.1 -37.2 -37.3 -37.4 -37.5 -37.6 -37.7 -37.8 -37.9 -38.0 -38.1 -38.2 -38.3 -38.4 -38.5 -38.6 -38.7 -38.8 -38.9 -39.0 -39.1 -39.2 -39.3 -39.4 -39.5 -39.6 -39.7 -39.8 -39.9 -40.0 -40.1 -40.2 -40.3 -40.4 -40.5 -40.6 -40.7 -40.8 -40.9 -41.0 -41.1 -41.2 -41.3 -41.4 -41.5 -41.6 -41.7 -41.8 -41.9 -42.0 -42.1 -42.2 -42.3 -42.4 -42.5 -42.6 -42.7 -42.8 -42.9 -43.0 -43.1 -43.2 -43.3 -43.4 -43.5 -43.6 -43.7 -43.8 -43.9 -44.0 -44.1 -44.2 -44.3 -44.4 -44.5 -44.6 -44.7 -44.8 -44.9 -45.0 -45.1 -45.2 -45.3 -45.4 -45.5 -45.6 -45.7 -45.8 -45.9 -46.0 -46.1 -46.2 -46.3 -46.4 -46.5 -46.6 -46.7 -46.8 -46.9 -47.0 -47.1 -47.2 -47.3 -47.4 -47.5 -47.6 -47.7 -47.8 -47.9 -48.0 -48.1 -48.2 -48.3 -48.4 -48.5 -48.6 -48.7 -48.8 -48.9 -49.0 -49.1 -49.2 -49.3 -49.4 -49.5 -49.6 -49.7 -49.8 -49.9 -50.0 -50.1 -50.2 -50.3 -50.4 -50.5 -50.6 -50.7 -50.8 -50.9 -51.0 -51.1 -51.2 -51.3 -51.4 -51.5 -51.6 -51.7 -51.8 -51.9 -52.0 -52.1 -52.2 -52.3 -52.4 -52.5 -52.6 -52.7 -52.8 -52.9 -53.0 -53.1 -53.2 -53.3 -53.4 -53.5 -53.6 -53.7 -53.8 -53.9 -54.0 -54.1 -54.2 -54.3 -54.4 -54.5 -54.6 -54.7 -54.8 -54.9 -55.0 -55.1 -55.2 -55.3 -55.4 -55.5 -55.6 -55.7 -55.8 -55.9 -56.0 -56.1 -56.2 -56.3 -56.4 -56.5 -56.6 -56.7 -56.8 -56.9 -57.0 -57.1 -57.2 -57.3 -57.4 -57.5 -57.6 -57.7 -57.8 -57.9 -58.0 -58.1 -58.2 -58.3 -58.4 -58.5 -58.6 -58.7 -58.8 -58.9 -59.0 -59.1 -59.2 -59.3 -59.4 -59.5 -59.6 -59.7 -59.8 -59.9 -60.0 -60.1 -60.2 -60.3 -60.4 -60.5 -60.6 -60.7 -60.8 -60.9 -61.0 -61.1 -61.2 -61.3 -61.4 -61.5 -61.6 -61.7 -61.8 -61.9 -62.0 -62.1 -62.2 -62.3 -62.4 -62.5 -62.6 -62.7 -62.8 -62.9 -63.0 -63.1 -63.2 -63.3 -63.4 -63.5 -63.6 -63.7 -63.8 -63.9 -64.0 -64.1 -64.2 -64.3 -64.4 -64.5 -64.6 -64.7 -64.8 -64.9 -65.0 -65.1 -65.2 -65.3 -65.4 -65.5 -65.6 -65.7 -65.8 -65.9 -66.0 -66.1 -66.2 -66.3 -66.4 -66.5 -66.6 -66.7 -66.8 -66.9 -67.0 -67.1 -67.2 -67.3 -67.4 -67.5 -67.6 -67.7 -67.8 -67.9 -68.0 -68.1 -68.2 -68.3 -68.4 -68.5 -68.6 -68.7 -68.8 -68.9 -69.0 -69.1 -69.2 -69.3 -69.4 -69.5 -69.6 -69.7 -69.8 -69.9 -70.0 -70.1 -70.2 -70.3 -70.4 -70.5 -70.6 -70.7 -70.8 -70.9 -71.0 -71.1 -71.2 -71.3 -71.4 -71.5 -71.6 -71.7 -71.8 -71.9 -72.0 -72.1 -72.2 -72.3 -72.4 -72.5 -72.6 -72.7 -72.8 -72.9 -73.0 -73.1 -73.2 -73.3 -73.4 -73.5 -73.6 -73.7 -73.8 -73.9 -74.0 -74.1 -74.2 -74.3 -74.4 -74.5 -74.6 -74.7 -74.8 -74.9 -75.0 -75.1 -75.2 -75.3 -75.4 -75.5 -75.6 -75.7 -75.8 -75.9 -76.0 -76.1 -76.2 -76.3 -76.4 -76.5 -76.6 -76.7 -76.8 -76.9 -77.0 -77.1 -77.2 -77.3 -77.4 -77.5 -77.6 -77.7 -77.8 -77.9 -78.0 -78.1 -78.2 -78.3 -78.4 -78.5 -78.6 -78.7 -78.8 -78.9 -79.0 -79.1 -79.2 -79.3 -79.4 -79.5 -79.6 -79.7 -79.8 -79.9 -80.0 -80.1 -80.2 -80.3 -80.4 -80.5 -80.6 -80.7 -80.8 -80.9 -81.0 -81.1 -81.2 -81.3 -81.4 -81.5 -81.6 -81.7 -81.8 -81.9 -82.0 -82.1 -82.2 -82.3 -82.4 -82.5 -82.6 -82.7 -82.8 -82.9 -83.0 -83.1 -83.2 -83.3 -83.4 -83.5 -83.6 -83.7 -83.8 -83.9 -84.0 -84.1 -84.2 -84.3 -84.4 -84.5 -84.6 -84.7 -84.8 -84.9 -85.0 -85.1 -85.2 -85.3 -85.4 -85.5 -85.6 -85.7 -85.8 -85.9 -86.0 -86.1 -86.2 -86.3 -86.4 -86.5 -86.6 -86.7 -86.8 -86.9 -87.0 -87.1 -87.2 -87.3 -87.4 -87.5 -87.6 -87.7 -87.8 -87.9 -88.0 -88.1 -88.2 -88.3 -88.4 -88.5 -88.6 -88.7 -88.8 -88.9 -89.0 -89.1 -89.2 -89.3 -89.4 -89.5 -89.6 -89.7 -89.8 -89.9 -90.0 -90.1 -90.2 -90.3 -90.4 -90.5 -90.6 -90.7 -90.8 -90.9 -91.0 -91.1 -91.2 -91.3 -91.4 -91.5 -91.6 -91.7 -91.8 -91.9 -92.0 -92.1 -92.2 -92.3 -92.4 -92.5 -92.6 -92.7 -92.8 -92.9 -93.0 -93.1 -93.2 -93.3 -93.4 -93.5 -93.6 -93.7 -93.8 -93.9 -94.0 -94.1 -94.2 -94.3 -94.4 -94.5 -94.6 -94.7 -94.8 -94.9 -95.0 -95.1 -95.2 -95.3 -95.4 -95.5 -95.6 -95.7 -95.8 -95.9 -96.0 -96.1 -96.2 -96.3 -96.4 -96.5 -96.6 -96.7 -96.8 -96.9 -97.0 -97.1 -97.2 -97.3 -97.4 -97.5 -97.6 -97.7 -97.8 -97.9 -98.0 -98.1 -98.2 -98.3 -98.4 -98.5 -98.6 -98.7 -98.8 -98.9 -99.0 -99.1 -99.2 -99.3 -99.4 -99.5 -99.6 -99.7 -99.8 -99.9 -100.0 -100.1 -100.2 -100.3 -100.4 -100.5 -100.6 -100.7 -100.8 -100.9 -101.0 -101.1 -101.2 -101.3 -101.4 -101.5 -101.6 -101.7 -101.8 -101.9 -102.0 -102.1 -102.2 -102.3 -102.4 -102.5 -102.6 -102.7 -102.8 -102.9 -103.0 -103.1 -103.2 -103.3 -103.4 -103.5 -103.6 -103.7 -103.8 -103.9 -104.0 -104.1 -104.2 -104.3 -104.4 -104.5 -104.6 -104.7 -104.8 -104.9 -105.0 -105.1 -105.2 -105.3 -105.4 -105.5 -105.6 -105.7 -105.8 -105.9 -106.0 -106.1 -106.2 -106.3 -106.4 -106.5 -106.6 -106.7 -106.8 -106.9 -107.0 -107.1 -107.2 -107.3 -107.4 -107.5 -107.6 -107.7 -107.8 -107.9 -108.0 -108.1 -108.2 -108.3 -108.4 -108.5 -108.6 -108.7 -108.8 -108.9 -109.0 -109.1 -109.2 -109.3 -109.4 -109.5 -109.6 -109.7 -109.8 -109.9 -110.0 -110.1 -110.2 -110.3 -110.4 -110.5 -110.6 -110.7 -110.8 -110.9 -111.0 -111.1 -111.2 -111.3 -111.4 -111.5 -111.6 -111.7 -111.8 -111.9 -112.0 -112.1 -112.2 -112.3 -112.4 -112.5 -112.6 -112.7 -112.8 -112.9 -113.0 -113.1 -113.2 -113.3 -113.4 -113.5 -113.6 -113.7 -113.8 -113.9 -114.0 -114.1 -114.2 -114.3 -114.4 -114.5 -114.6 -114.7 -114.8 -114.9 -115.0 -115.1 -115.2 -115.3 -115.4 -115.5 -115.6 -115.7 -115.8 -115.9 -116.0 -116.1 -116.2 -116.3 -116.4 -116.5 -116.6 -116.7 -116.8 -116.9 -117.0 -117.1 -117.2 -117.3 -117.4 -117.5 -117.6 -117.7 -117.8 -117.9 -118.0 -118.1 -118.2 -118.3 -118.4 -118.5 -118.6 -118.7 -118.8 -118.9 -119.0 -119.1 -119.2 -119.3 -119.4 -119.5 -119.6 -119.7 -119.8 -119.9 -120.0 -120.1 -120.2 -120.3 -120.4 -120.5 -120.6 -120.7 -120.8 -120.9 -121.0 -121.1 -121.2 -121.3 -121.4 -121.5 -121.6 -121.7 -121.8 -121.9 -122.0 -122.1 -122.2 -122.3 -122.4 -122.5 -122.6 -122.7 -122.8 -122.9 -123.0 -123.1 -123.2 -123.3 -123.4 -123.5 -123.6 -123.7 -123.8 -123.9 -124.0 -124.1 -124.2 -124.3 -124.4 -124.5 -124.6 -124.7 -124.8 -124.9 -125.0 -125.1 -125.2 -125.3 -125.4 -125.5 -125.6 -125.7 -125.8 -125.9 -126.0 -126.1 -126.2 -126.3 -126.4 -126.5 -126.6 -126.7 -126.8 -126.9 -127.0 -127.1 -127.2 -127.3 -127.4 -127.5 -127.6 -127.7 -127.8 -127.9 -128.0 -128.1 -128.2 -128.3 -128.4 -128.5 -128.6 -128.7 -128.8 -128.9 -129.0 -129.1 -129.2 -129.3 -129.4 -129.5 -129.6 -129.7 -129.8 -129.9 -130.0 -130.1 -130.2 -130.3 -130.4 -130.5 -130.6 -130.7 -130.8 -130.9 -131.0 -131.1 -131.2 -131.3 -131.4 -131.5 -131.6 -131.7 -131.8 -131.9 -132.0 -132.1 -132.2 -132.3 -132.4 -132.5 -132.6 -132.7 -132.8 -132.9 -133.0 -133.1 -133.2 -133.3 -133.4 -133.5 -133.6 -133.7 -133.8 -133.9 -134.0 -134.1 -134.2 -134.3 -134.4 -134.5 -134.6 -134.7 -134.8 -134.9 -135.0 -135.1 -135.2 -135.3 -135.4 -135.5 -135.6 -135.7 -135.8 -135.9 -136.0 -136.1 -136.2 -136.3 -136.4 -136.5 -136.6 -136.7 -136.8 -136.9 -137.0 -137.1 -137.2 -137.3 -137.4 -137.5 -137.6 -137.7 -137.8 -137.9 -138.0 -138.1 -138.2 -138.3 -138.4 -138.5 -138.6 -138.7 -138.8 -138.9 -139.0 -139.1 -139.2 -139.3 -139.4 -139.5 -139.6 -139.7 -139.8 -139.9 -140.0 -140.1 -140.2 -140.3 -140.4 -140.5 -140.6 -140.7 -140.8 -140.9 -141.0 -141.1 -141.2 -141.3 -141.4 -141.5 -141.6 -141.7 -141.8 -141.9 -142.0 -142.1 -142.2 -142.3 -142.4 -142.5 -142.6 -142.7 -142.8 -142.9 -143.0 -143.1 -143.2 -143.3 -143.4 -143.5 -143.6 -143.7 -143.8 -143.9 -144.0 -144.1 -144.2 -144.3 -144.4 -144.5 -144.6 -144.7 -144.8 -144.9 -145.0 -145.1 -145.2 -145.3 -145.4 -145.5 -145.6 -145.7 -145.8 -145.9 -146.0

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP DEG C
0415	35 16.9	123 17.3	489.1	2.29	0.79	2.9			15.3
			489.7	0.0	0.80	0.0			15.3
			489.3	0.0	0.81	0.0			15.3
			490.0	0.0	0.61	0.0			15.3
			490.6						15.3
			491.2						15.3
			491.9						15.3
			492.5						15.3
			493.1						15.3
			493.8						15.2
			494.4						15.1
			495.0						15.1
			495.7						15.1
			496.3						15.1
			496.9						15.1
			497.6						15.1
0505	35 22.7	123 11.7	498.2						15.0
			498.8						15.0
			499.5						15.0
			500.1						15.0
			500.7						15.0
			501.4						15.0
			502.0						15.0
			502.6						15.0
			503.3						15.0
			503.9						15.0
			504.5						15.0
			505.2						15.0
			505.8						15.0
			506.4						15.0
			507.1						15.0
			507.7						15.1
			508.3						15.1
			509.0						15.1
			509.6						15.1
			510.2						15.1
			510.9						15.1
			511.5						15.0
			512.1						15.0
			512.8						15.1
			513.4						15.3
			514.1						15.3
			514.7						15.3
			515.3						15.3
			516.0						15.3
			516.6						15.3
			517.2						15.3
			517.9						15.3
			518.5						15.4
			519.1						15.4

R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE(CRUIS= VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR.RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP DEG C
0600	35	28.6	123	6.2					
			519.8						15.4
			520.4						15.4
			521.0						15.4
			521.7						15.4
			522.3						15.4
			523.0						15.4
			523.6						15.4
			524.2						15.4
			524.8						15.4
			525.5						15.4
			526.1						15.4
			526.7						15.4
			527.4						15.4
			528.0						15.4
			528.6						15.4
			529.3						15.4
			529.9						15.4
			530.5						15.4
			531.2						15.4
			531.8						15.4
			532.4						15.4
			533.1						15.4
			533.7						15.4
			534.3						15.4
			535.0						15.4
			535.6						15.4
			536.2						15.4
			536.9						15.4
			537.5						15.4
			538.1						15.4
			538.8						15.4
			539.4						15.4
			540.0						15.4
			540.7						15.4
			541.3						15.4
			541.9						15.4
			542.6						15.4
			543.2						15.4
			543.8						15.4
			544.5						15.4
			545.1						15.4
			545.7						15.4
			546.4						15.4
			547.0						15.4
			547.6						15.4
			548.3						15.4
			548.9						15.4
			549.5						15.4
			550.2						15.4
			550.8						15.4

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR.RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP DEG C
0900	35 49.2	122 46.6	583.1						15.4
			584.8						15.4
			585.0						15.4
			585.7						15.3
			586.3						15.3
			587.6						15.3
			588.2						15.3
			589.5						15.3
			590.1						15.3
			590.7						15.3
			591.4						15.3
			592.0						15.3
			593.3						15.3
			593.9						15.4
			594.5						15.4
1000	35 56.5	122 38.9	595.2						15.4
			596.8						15.4
			597.4						15.4
			598.3						15.4
			599.0						15.4
			600.2						15.4
			600.9						15.4
			601.5						15.4
			602.1						15.4
			602.8						15.4
			603.4						15.4
			604.0						15.3
			604.7						15.3
			605.3						15.3
			606.0						15.0
			606.6						14.8
			607.2						14.6
			607.9						14.6
			608.5						14.5
			609.1						14.5
			609.8						14.5
			610.4						14.4
			611.0						14.4
			611.7						14.4
			612.3						14.4
			612.9						14.4
			613.6						14.3
			614.2						14.3

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP DEC C
1100	36	3.1	122 32.2						
			614.8						14.3
			615.5						14.3
			616.1						14.3
			616.7						14.3
			617.4						14.3
			618.0						14.3
			619.6						14.3
			619.3						14.3
			619.9						14.3
			620.5						14.3
			621.2						14.3
			621.8						14.3
			622.4						14.3
			623.1						14.3
			624.3						14.3
			625.0						14.3
			625.6						14.3
			626.2						14.3
			626.9						14.3
			627.5						14.3
			628.1						14.3
			628.9						14.3
			629.4						14.3
			630.0						14.3
			630.7						14.3
			631.3						14.3
			631.9						14.3
			632.6						14.3
			633.2						14.3
			633.8						14.3
			634.5						14.3
			635.1						14.3
			635.7						14.3
			636.4						14.3
			637.0						14.3
			637.6						14.3
			638.3						14.3
			639.9						14.3
			640.5						14.3
			640.8						14.3
			641.4						14.3
			642.1						14.3
			643.3						14.3
			644.0						14.3
			644.6						14.3
			645.2						14.3
			645.9						14.3

APPENDIX B

STRESS MAG
DE JON =
5 DMS PER 50 CM

EKMAN TRAN
DE JON =
5 DMS PER 50 CM


UP INDEX
DE JON =
5 DMS PER 50 CM

VERT VELOC
DE JON =
5 DMS PER 50 CM


1 MAR 1979
2 MAR 1979
3 MAR 1979
4 MAR 1979
5 MAR 1979
6 MAR 1979
7 MAR 1979
8 MAR 1979
9 MAR 1979
10 MAR 1979
11 MAR 1979
12 MAR 1979
13 MAR 1979
14 MAR 1979
15 MAR 1979
16 MAR 1979
17 MAR 1979

19 MAR 1979
20 MAR 1979
21 MAR 1979
22 MAR 1979
23 MAR 1979
24 MAR 1979
25 MAR 1979
26 MAR 1979
27 MAR 1979
28 MAR 1979
29 MAR 1979
30 MAR 1979

N



31 MAR 1979
 1 APR 1979
 2 APR 1979
 3 APR 1979
 4 APR 1979
 5 APR 1979
 6 APR 1979
 7 APR 1979
 8 APR 1979
 9 APR 1979
 10 APR 1979
 11 APR 1979
 12 APR 1979
 13 APR 1979
 14 APR 1979
 15 APR 1979
 16 APR 1979
 17 APR 1979
 18 APR 1979
 19 APR 1979
 20 APR 1979
 21 APR 1979
 22 APR 1979
 23 APR 1979
 24 APR 1979
 25 APR 1979
 26 APR 1979
 27 APR 1979
 28 APR 1979
 29 APR 1979
 30 APR 1979
 1 MAY 1979
 2 MAY 1979



STRESS MRG

DE 1901 =
5 OBS / SEC

EKMAN TRAN

DE 1901 =
5 OBS / SEC

UP INDEX

DE 1901 =
5 OBS / SEC

VERT VELOC

DE 1901 =
1 M / SEC

1 JUN 1979

2 JUN 1979

3 JUN 1979

4 JUN 1979

5 JUN 1979

6 JUN 1979

7 JUN 1979

8 JUN 1979

9 JUN 1979

10 JUN 1979

11 JUN 1979

12 JUN 1979

13 JUN 1979

14 JUN 1979

15 JUN 1979

16 JUN 1979

17 JUN 1979

18 JUN 1979

19 JUN 1979

20 JUN 1979

21 JUN 1979

22 JUN 1979

23 JUN 1979

24 JUN 1979

25 JUN 1979


26 JUN 1979

27 JUN 1979

28 JUN 1979

29 JUN 1979

30 JUN 1979





BIBLIOGRAPHY

1. Anonymous, Technicon Industrial Systems Technical Publication No. TA 1-0170-20, Operation Manual for the Technicon Auto Analyzer II System, 1972.
2. Anonymous, Technicon Industrial Systems Industrial Method No. 177-72-WM, Ortho Phosphate in Water and Seawater, January 1973.
3. Anonymous, Technicon Industrial Systems Industrial Method No. 175-72-WM, Nitrate and Nitrite in Water and Seawater, January 1973.
4. Anonymous, Technicon Industrial Systems Industrial Method No. 100-70-W/B, Nitrate and Nitrite in Water and Waste Water, January 1978.
5. Bakun, A., NOAA Technical Report NMFS SSRF-671, Coastal Upwelling Indices, West Coast of North America 1946-71, 1973.
6. Banse, K., "The Nitrogen-to-Phosphorus Ratio in the Photic Zone of the Sea and the Elemental Composition of the Plankton," Deep Sea Research, V. 21, pp. 767-771.
7. Barton, E. D., Huyer, A. and Smith, R. L., "Temporal Variations Observed in the Hydrographic Regime Near Cabo Corbeiro in the Northwest African Upwelling Region February to April 1974," Deep Sea Research, V. 24, pp. 7-25, 1977.
- *8. Broecker, W. S., Chemical Oceanography, Harcourt Brace Jovanovich, pp. 9-13, 1974.
9. Bower, R. L., Gohrband, H. S., Pichel, W. G., Signore, T. L., and Walter, C. C., NOAA Technical Memorandum NESS 78, Satellite Derived Sea Surface Temperatures from NOAA Spacecraft, pp. 1-12 and 35-39, June 1976.
10. Butler, E. I., Knox, S., and Liddicott, M. I., "The Relationship Between Inorganic and Organic Nutrients in Seawater," Journal of the Marine Biological Association of the United Kingdom, V. 59, pp. 239-250, 1979.
11. Cheney, R. E., and Winfrey, D. E., NAVOCEANO Technical Note 3700-56-76, Distribution and Classification of Fronts, p. 206, August 1976.

*Background literature not cited in the text.

12. Codispoti, L. A., and Friederich, G. E., "Local and Mesoscale Influences on Nutrient Variability in the Northwest African Upwelling Region Near Cabo Corbeiro," Deep Sea Research, V. 25, pp. 751-770, 1977.
13. Codispoti, L. A., Dugdale, R. C., Minas, H. J., "A Comparison of the Nutrient Regimes Off Northwest Africa, Peru and Baja California," CUEA Newsletter, V. 8, pp. 2-25, January 1979.
14. Dugdale, R. C., "Nutrient Limitation in the Sea: Dynamics, Identification, and Significance," Limnology and Oceanography, V. 12, pp. 685-695, 1967.
15. Dugdale, R. C., and Georing, J. J., "Uptake of New and Regenerated Forms of Nitrogen in Primary Productivity," Limnology and Oceanography, V. 12, pp. 196-206, 1962.
16. Fairbridge, R. W., The Encyclopedia of Oceanography, Reinhold Publishing Corporation, 1966.
17. Fett, R. W., La Violetter, P. E., Nestor, M., Nickerson, J. W., Rabe, K., NAVENVPREDRSHFAC Technical Report No. 77-04, Navy Tactical Applications Guide, Vol. II, Environmental Phenomena and Effects, pp. 3B-6, January 1979.
18. Holm-Hansen, O. and Karl, D. M., "Effects of Luciferin Concentration on the Quantitative Assay of ATP Using Crude Luciferase Preparations," Analytical Biochemistry, V. 75, pp. 100-112, 1976.
19. Holm-Hansen, O., "Determination of Microbial Biomass in Ocean Profiles," Limnology and Oceanography, V. 14, pp. 740-747, 1969.
20. Huyer, A., Smith, R. L. and Pillsburg, R. D., "Observations in a Coastal Upwelling Region During a Period of Variable Winds (Oregon Coast, July 1972)," Tethys, V. 6, pp. 1-2, 1974.
21. Huntsman, S. A. and Barber, R. T., "Primary Production Off Northwest Africa the Relationship to Wind and Nutrient Conditions," Deep Sea Research, V. 24, pp. 25-35, 1977.
22. Jones, P. G. W., "The Southern Benguela Current Region in February 1966: Part 1 Chemical Observations with Particular Reference to Upwelling," Deep Sea Research, V. 18, pp. 193-208, 1971.
23. Kelly, J. C., Whitledge, T. E. and Dugdale, R. C., "Results of Sea Surface Mapping in the Peru Upwelling System," Limnology and Oceanography, V. 20, pp. 784-794, 1975.

24. Legeckis, R. V., "A Survey of Worldwide Sea Surface Temperature Fronts Detected by Environmental Satellites," Journal of Geophysical Research, V. 83, pp. 4501-4523, 20 September 1978.
25. Lorenzen, C. J., "A Method for the Continuous Measurement of In Vivo Chlorophyll Concentration," Deep Sea Research, V. 13, pp. 223-227, 1966.
26. Paulson, G. P., A Study of Nutrient Variations in the Surface and Mixed Layer of Monterey Bay Using Automatic Analysis Techniques, Master's Thesis, Naval Postgraduate School, Monterey, California, September 1972.
27. Redfield, A. C., "On the Proportions of Organic Derivatives in Seawater and Their Relation to the Composition of Plankton," James Johnstone Memorial Volume, University Press of Liverpool, pp. 176-192, 1934.
28. Redfield, A. C., "The Biological Control of Chemical Factors in the Environment," American Scientist, pp. 205-221, September 1958.
29. Redfield, A. C., Ketchum, B. H., and Richards, F. A., "The Influence of Organisms on the Composition of Seawater," in N. M. Hill (ed.), The Sea, V. 2, pp. 26-27, Interscience Publishers, 1963.
30. Reid, J. L., Roden, G. I., and Wyllie, J. G., "Studies of the California Currents Systems," California Cooperative Oceanic Fisheries Investigation, Progress Report, pp. 27-56, 1 July 1956-1 January 1958.
31. Richards, F. A., "Anoxic Basins and Fjords," in J. P. Riley and G. Skirrow (ed.), Chemical Oceanography, V. 1, Academic Press, pp. 611-645, 1965.
32. Riley, J. P., and Skirrow, G., Chemical Oceanography, 2nd Ed., V. 3, Academic Press, 1975.
33. SCOR Working Group 36 (with ACMRR and ACOMR), Coastal Upwelling Processes, Report of 2nd Meeting, Kiel, FRG 24-28 June 1974, CUEA Newsletter, V. 4, pp. 12-20, September 1975.
34. Shaffer, G., "On Quasi-Steady Three-Dimensional Coastal Upwelling Circulation," CUEA Newsletter, V. 5, pp. 25-29, March 1976.
- *35. Sherman, J. W., III, "Current and Future Satellites for Ocean Monitoring," In: Proceedings of the Eleventh International Symposium on Remote Sensing of Environment, Environmental Research Institute of Michigan, Ann Arbor, Michigan, V. 1, pp. 288-299, 1977.

36. Smith, S. L., and Whittledge, T. E., "The Role of Zooplankton in the Regeneration of Nitrogen in a Coastal Upwelling System Off Northwest Africa," Deep Sea Research, V. 24, pp. 49-56, 1977.
37. Strickland, J. D. H. and Parsons, T. R., A Practical Handbook of Seawater Analysis, J. C. Stevenson (Ed.), Queen's Printer and Comptroller of Stationery, pp. 49-62 and 71-76, 1968.
38. Sverdrup, H. U., Johnson, M. W., and Fleming, R. H., The Oceans, Their Physics, Chemistry, and General Biology, Prentice Hall, 1942.
39. Thomas, W. H., and Siebert, D. L. R., "Distribtuion of Nitrate, Nitrite, Phosphate, and Silicate in the California Current Region, 1969," California Cooperative Oceanic Fisheries Investigations, V. 20, June 1974.
40. Thomas, W. H., "Phytoplankton Nutrient Enrichment Experiments Off Baja California and in the Eastern Equatorial Pacific Ocean," Journal of the Fisheries Research Board of Canada, V. 26, pp. 1133-1145, 1969.
41. Traganza, E. D., Chemical Mesoscale and ATP-Biomass Correlated with Satellite Imagery of Oceanic Fronts, Interim Report on Proposal Submitted to Office of Naval Research, Code 482, 27 June 1978.
42. Traganza, E. D., Naval Postgraduate School Technical Report No. 68-79-008, "The Use of Temperature and Color in Satellite Detection of Ocean Fronts and Eddies for ASW Applications," February 1979.
43. Traganza, E. D., Nestor, D. A., and McDonald, A. K., "Satellite Observations of a Nutrient Upwelling Off the Coast of California," (in press Journal of Geophysical Research), 1980.
44. Treguer, P., and LeCorre, P., "The Ratios of Nitrate Phosphate, and Silicate During Uptake and Regeneration Phases of the Moroccan Upwelling Regime," Deep Sea Research, V. 23A, pp. 163-184, 1977.
- *45. Wiebe, P. H., "Gulf Stream Cold Core Rings: Large Scale Interaction Sites for Open Ocean Plankton Communities," Deep Sea Research, V. 23, pp. 695-710, 1978.

INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	2
2. Library, Code 0142 Naval Postgraduate School Monterey, California 93940	2
3. Chairman, Code 68 Department of Oceanography Naval Postgraduate School Monterey, California 93940	3
4. Director Naval Oceanography Division (OP952) Navy Department Washington, DC 20350	1
5. Office of Naval Research Code 480 Naval Ocean Research and Development Activity NSTL Station, Mississippi 39529	1
6. Dr. Robert E. Stevenson Scientific Liaison Office, ONR Scripps Institution of Oceanography La Jolla, California 92037	1
7. SIO Library University of California, San Diego P.O. Box 2367 La Jolla, California 92037	1
8. Department of Oceanography Library University of Washington Seattle, Washington 98105	1
9. Department of Oceanography Library Oregon State University Corvallis, Oregon 97331	1
10. Commanding Officer Fleet Numerical Weather Central Monterey, California 93940	1

11. Commanding Officer 1
Naval Environmental Prediction Research
Facility
Monterey, California 93940
12. Commander 1
Oceanographic Systems Pacific
Box 1390
Pearl Harbor, Hawaii 96860
13. Commanding Officer 1
Naval Ocean Research and Development
Activity
Dr. Edward Green
Code 482
NSTL Station, Mississippi 39529
14. Commander 1
Naval Oceanography Command
NSTL Station, Mississippi 39529
15. Commanding Officer 1
Naval Oceanographic Office
NSTL Station, Mississippi 39529
16. Dr. E. D. Traganza, Code 68Tg 5
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940
17. Mr. Jerry Norton, Code 68 1
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940
18. LT Don A. Nestor 1
Helicopter Antisubmarine Squadron One
Naval Air Station
Jacksonville, Florida 32212
19. Mr. Laurence Breaker 1
National Environmental Satellite Service
660 Price Avenue
Redwood City, California 94063
20. Mr. Roland Nagle 1
Naval Environmental Prediction Research
Facility
Monterey, California 93940
21. Mr. Morton Kronengold 1
Institute for Acoustical Research
615 SW 2nd Avenue
Miami, Florida 33130

22. Commander 3
Area ASW Forces Sixth Fleet
LCDR John W. Conrad
U.S. Naval Support Activity, Naples
FPO New York 09521
23. Mr. Rolf J. Berger, Code 68 1
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940
24. LT Walter Hanson, Code 35 1
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940
25. Ms. Andrea McDonald, Code 61 1
Department of Chemistry
Naval Postgraduate School
Monterey, California 93940
26. Captain W. W. Reynolds, Code 35 1
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940
27. Dr. Stevens P. Tucker, Code 68Tx 1
Department of Oceanography
Naval Postgraduate School
Monterey, California 93940

Thesis
C7415
c.1

Conrad

189005

Relationships between
sea surface temperature
and nutrients in satel-
lite detected oceanic
fronts.

Th
C7
c.

Thesis
C7415
c.1

Conrad

189005

Relationships between
sea surface temperature
and nutrients in satel-
lite detected oceanic
fronts.

thesC7415

Relationships between sea surface temper



3 2768 002 09345 2

DUDLEY KNOX LIBRARY